



Perspectives

R. B. Woodward's unfinished symphony: designing organic superconductors (1975–79)[☆]

Michael P. Cava^{a,†}, M.V. Lakshmikantham^{a,†}, Roald Hoffmann^b, Robert M. Williams^{c,*}

^a Department of Chemistry, The University of Alabama, Tuscaloosa, AL 35487, USA

^b Department of Chemistry and Chemical Biology, Cornell University, Ithaca, NY 14853, USA

^c Department of Chemistry, Colorado State University, Fort Collins, CO 80523, USA

ARTICLE INFO

Article history:

Received 27 February 2011

Accepted 3 May 2011

Available online 12 May 2011

Keywords:

Organic conducting materials

Polymers

Sulfur-nitrogen polymers

History of chemistry

Contents

Introduction	6771
Preface	6772
Robert Burns Woodward—an architect of desire at his drawing desk	6772
Woodward's Thoughts, Drawings, and Work on Potential Organic Superconductors	6773
1. Modifications of (SN) _x	6773
Acknowledgements	6796
References and notes	6796
Biographical sketch	6797

Introduction

When Robert Burns Woodward passed away in July 1979, he left behind 699 pages of hand-written notes in a neat stack in his top desk drawer. These notes were kept safely by his youngest son Eric Woodward in a small cardboard box for many years while the family pondered what to do with them. In 2001, Woodward's two grand daughters digitally scanned each page and numbered them in the order in which they were left in the original stack to protect the images for archival purposes. The notes were written on all

different sorts of paper, including hotel stationary, yellowed paper pads, white photocopy paper and light blue paper. These notes revealed that RBW was deeply immersed in an astonishing project that had consumed him during the last years of his life. He became passionately interested in designing, de novo, molecules that would be organic superconducting materials.

From interviews with the Woodward family, and some colleagues, it was apparent that RBW had difficulty convincing co-workers at Harvard, as well as at the Woodward Research Institute in Basel, Switzerland, to engage in much experimental work on the ideas he was formulating. Nonetheless, RBW appeared to press on thinking deeply about unique new structures that might have the capacity to efficiently conduct electricity.

After extensive discussions with the Woodward family, permission was secured to identify a close colleague or former co-worker of RBW, with expertise in this area, to examine these notes carefully and write a retrospective view of this last, unfinished

[☆] The actual scans of R. B. Woodward's notes are property of Eric and Crystal Woodward and shall not be published nor reproduced for commercial purposes without their permission.

* Corresponding author. E-mail address: rmw@lamar.colostate.edu (R.M. Williams).

† Deceased.

work of RBW. Prof. Michael P. Cava, a former graduate student of Woodward's at Harvard, stepped forward to tackle this assignment. The accompanying paper, which primarily covers structures with, which Prof. Cava was most comfortable commenting on, is but a small sampling of the myriad ideas Woodward put on paper. I hope that the complete set of these notes will one day reach the chemical community; it is of significant historical significance.

Woodward was best known of course, for his published work on natural products structure determination and their total synthesis. What is stunning about these notes is the mathematical and geometrical depth with, which Woodward was conceptualizing conducting materials. Roald Hoffmann interacted with Woodward from time-to-time on this project over the years and Hoffmann's Preface to this article adds significant depth, color, and perspective. Only two papers were published by Woodward in this area; the vast bulk of his ideas in this field, were deprived of ever having been experimentally tested. One can not help wondering had Woodward lived longer, what advances he might have pioneered in this field. This article, and the accompanying scanned images from his notes, give an amazing glimpse of Woodward's thought process, creativity, and deep intellectual gifts that characterized his brilliant career.

Robert M. Williams
Colorado State University

Preface

Robert Burns Woodward—an architect of desire at his drawing desk

In the last three years of his life, R. B. Woodward was gripped by the idea of designing and synthesizing an organic superconductor. The evidence to that creative obsession (and I use the word in its most positive sense) is to be found in the hundreds of meticulous drawings of molecules he left behind at his untimely death in 1979. These drawings, and the associated geometrical and algebraic ruminations, effectively comprise a sketch book, tracing the creative path of a great scientist impelled by molecular desire. In an architectural way, Woodward's blue and yellow pages represent the working through of the design of electronic function. By a master builder of molecules. Ergo the architect, and his drawing desk.

Organic conductors were hot in the mid-1970s. As yet they were a negative contribution to the gross domestic product of the countries where they were studied. No one had yet made an organic superconductor of any transition temperature of note. Buckminsterfullerene, not to speak of its salts, was just a theoretician's dream.

Yet Woodward was moved by that dream, and seized on the curious $(SN)_x$ system, as inorganic as it was organic, and on two-dimensional all-carbon lattices as starting points in his intellectual game. In the molecules he designed you see the workings of the organic imagination at its best, creative flights of molecular fancy. For example, if $(SN)_x$ did not have a handle on it, then the isoelectronic $(SCR)_x$ would. And what an incredible variety of structures surface once this ingenious molecular architect set his mind to it!

The properties desired were conductivity and superconductivity. And now Woodward was faced with a problem—not chemists, but theoretical physicists shaped the language and the ways of thinking about conductivity. Band structures, electron mobilities, scattering off phonons, Fermi surfaces were not part of the language of chemists.

Woodward found a model a chemist could wrap his mind around, Little's graphic concept of a chain of radicals surrounded by polarizable material, and the availability of vibrations decreasing electron repulsion. RBW complemented this picture by the drawing of resonance structures—in these pages we see him tirelessly redrawing the molecules to display the resonance structures rippling across them.

We don't see many molecular orbitals; this is interesting. For the facility RBW acquired in orbital thinking, beginning with the ferrocene story and the octant rule, was reinforced by the interaction he and I had in the orbital symmetry control story. In that marvelous collaboration, I was transformed from a calculator of MOs to a builder of chemical explanations based on them and their interactions.

But there were so many MOs in 1-D, 2-D, and 3-D extended structures, the real and hypothetical polymers adorning Woodward's pages. A single molecular orbital at the Fermi level did not have the 'power' to influence anything. One had to find a way of thinking about bunches of orbitals, a language that bridged solid state physics thinking with frontier orbital arguments. And at first RBW did not command that language. He drew me into the problem by asking questions. For the first time in seven years (the long *Angewandte Chemie* account of our work was published in 1969) we worked together. A brilliant postdoc of mine from Korea via Canada, Myung-Hwan Whangbo, was drawn to the project. He wrote our first extended Hückel theory band program. And he, I, and RBW began to try to understand bands in a chemical way.

For me, it was the beginning of a path I am still on—from bands to bonds and back again, shaping understanding at the interface of chemistry and physics.

One thing you see at intervals in the Woodward organic conductor pages is a working out of algebraic sums—favorable electron counts, interlayer spacings, the geometric constraints in the fitting in a molecule into a polygon. At various times in his life, RBW said that he flirted with mathematics when he was young. That if he had not become a chemist, he would have been a mathematician. His old love, his fearlessness—no, delight—in mathematical complication, is evident in these pages.

And the drawings! I found one page in the set that is carelessly done. It's mine. Woodward's are precise, drawn with extreme care. In free-hand lines, firmly straight where they should be, in polygons, even shaded and colored in, as in his fillings of the plane with 24-membered $C_6S_6N_{12}$ rings, the architectonic imagination soars in these drawings.

In another context, it may not be surprising that the children of this molecular architect and his artist wife would become an artist and an architect.

Like many people, Woodward had trouble translating his dreams into reality. The stellar young people who gravitated from around the world toward his Cambridge lab, often with their own support, came (and had been sent there) to gain mastery in organic synthesis. Main-group and polymer chemistry that were needed to make the SCN polymers were not part of the skill set of these young acolytes.

In Basel, Ciba–Geigy had set up a laboratory for Woodward's research. The co-workers here were more senior researchers, and in principle should do whatever RBW wanted them to do. But even if Ciba–Geigy was committed to a free-ranging Woodward, organic conducting systems were too much for them. The Woodward Research Institute group made a pair of fascinating molecules mentioned in this paper, and published the work after RBW's death. But somehow the features of interest evident in the structures of these molecules did not make an impression on the team or their sponsors. The work was not followed up.

No point in thinking about the attractive paths that the chemistry of conducting materials could have taken, would have taken had Woodward lived. Let's be thankful for what is given to us in these crumbling yellow and blue pages. In them I see evidence of an inventive mind roaming widely, searching for an organic way to think about conductivity, designing in this microuniverse of chemistry singular molecules that are simply beautiful. And ... still waiting to be made.




Fig. 4. Polyacetylene and RBW notes, p 9.




Fig. 5. RBW notes, p 512.

Woodward and co-workers from Harvard, Ciba–Geigy and the Woodward Research Institute in Basel, the reaction of amidine with sulfur dichloride was shown to be very complex and no polymer was produced. However, the reaction of benzamidine (**18**) with sulfur dichloride gave a very low yield of an unusual eight-membered heterocycle **19**, the structure of which was confirmed by X-ray analysis⁸ (Fig. 10).

The eight-membered ring was practically flat, and all the S–N distances were equal as were the C–N bond lengths, indicative of a delocalized 10π -electron aromatic system. In contrast, the reaction of *N,N*-dimethylguanidine and sulfur dichloride furnished **20** (Fig. 11), analogous to **19**, although its crystal structure showed a puckering due to folding along a central S–S axis.

The UV–vis spectra of **19** and **20** were distinct. The photoelectron spectrum (PES) of **19** and **20** were determined subsequently by several groups. Ab initio calculations were also carried out and found to be in agreement with the experimental results from X-ray photoelectron spectroscopy. The basic parent ring system is aromatic and has a planar structure.⁹ However, structure **20** as well as the *tert*-butyl-substituted derivative, preferred a puckered geometry with a transannular S–S bond. The factors that influence the geometric preferences of this fascinating ring system, related to S_4N_4 , remain to be elucidated.




Fig. 6. Polythiazyl and RBW notes, p 1, 63, 83.




Fig. 7. RBW notes, p 64.

Improved synthetic methods have also been reported as well as the isolation of a new product **21** (Fig. 12) from the reaction of benzamidine or its N–S trimethylsilyl derivative with sulfur chloride. Indeed, **21** was the major product in this reaction.¹⁰

Some selenium analogs have also been reported.¹¹ Heterocycle **8** has been studied since 1980 from both the synthetic and theoretical points of view.¹² The N_4S molecule **9** has also been subjected to theoretical calculations.¹³

Benzobisthiadiazole **11** was first reported in 1994 and was subsequently the subject of theoretical studies.¹⁴ Several selenium analogs are also known. Some derivatives of **11** bearing electron-donating groups are highly colored, because of their low HOMO–LUMO gap.

Bisthiadiazine **12** was described in 1978.¹⁵ Subsequently a number of theoretical studies and experimental verifications via X-ray, PES, CV, ESY etc. were carried out.^{15,16} Thiadiazolo thiadiazole **13** has been known from 1975 and is a very stable molecule.¹⁷ It has attracted its share of attention from theoretical chemists.^{14c,18}

There exist two references to the sulfur–nitrogen heterocycle **22**. The first pertains to the calculation in 1988 of topological resonance energies of S–N heterocycles.¹⁹ The second is the synthesis of an analog, **23**, isoelectronic with **22** (Fig. 13).²⁰

The oligomeric/polymeric phenylene thiodimide is not known diphenylsulfur dimides however, have been described, and well characterized since 1961.²¹

Woodward's thoughts appear to have focused on the 24-membered heterocycle **24** (Fig. 14; yet unknown) and isoelectronic analogs. In the hypothetical molecule **24**, three sulfurs can be considered internal (S_i), as opposed to the three external ones (S), which fall on the vertices of an isosceles triangle. The S_i – S_i distance can be calculated easily using simple trigonometrical principles, and using standard $-\text{N}=\text{C}-\text{N}-$ bond lengths and bond angles; It is approximately 7.64 Å. The center point of this triangle is also the center point of the 24-membered heterocycle.

It was Woodward's concept (Fig. 15) that in order to force the formation of the 24-membered ring, guanidine could be used as a template, which utilizes the propensity of hydrogen-bonding between the ring N and the guanidine N–H bond.

Triaminoborane was also considered. However, it is an unknown molecule and all indications are that it is an unstable entity.


Fig. 8. Some ring structures for $(\text{SN})_x$.

Table 1
Some N–S structures from the RBW notes





















































Structure & Notes Page #	Original Image RBW Notes
 <p>(p. 184, 185)</p>	
 <p>(p. 187)</p>	
 <p>(p. 209)</p>	
 <p>(p. 210; 213)</p>	
 <p>(p. 260)</p>	
 <p>(p. 214)</p>	
 <p>(p. 185)</p>	
 <p>(p. 348)</p>	

Table 1 (continued)

Structure & Notes Page #	Original Image RBW Notes
 <p>(p. 351)</p>	 <p>This would have one extra electron and one net negative charge per ring • put metal in middle? $M^+ \rightarrow M^+ \leftrightarrow \text{ring}$ would it then be a polyradical and paramagnetic? check with repeating unit</p>
 <p>(p. 125; 355)</p>	
 <p>(p. 355)</p>	
 <p>(p.211; 601)</p>	 <p>HOTEL DREI KÖNIGE AM RHEIN BASEL HO-11 1175 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2722 2723 2724 2725 2726 2727 2728 2729 2730 2731 2732 2733 2734 2735 2736 2737 2738 2739 2740 2741 2742 2743 2744 2745 2746 2747 2748 2749 2750 2751 2752 2753 2754 2755 2756 2757 2758 2759 2760 2761 2762 2763 2764 2765 2766 2767 2768 2769 2770 2771 2772 2773 2774 2775 2776 2777 2778 2779 2780 2781 2782 2783 2784 2785 2786 2787 2788 2789 2790 2791 2792 2793 2794 2795 2796 2797 2798 2799 2800 2801 2802 2803 2804 2805 2806 2807 2808 2809 2810 2811 2812 2813 2814 2815 2816 2817 2818 2819 2820 2821 2822 2823 2824 2825 2826 2827 2828 2829 2830 2831 2832 2833 2834 2835 2836 2837 2838 2839 2840 2841 2842 2843 2844 2845 2846 2847 2848 2849 2850 2851 2852 2853 2854 2855 2856 2857 2858 2859 2860 2861 2862 2863 2864 2865 2866 2867 2868 2869 2870 2871 2872 2873 2874 2875 2876 2877 2878 2879 2880 2881 2882 2883 2884 2885 2886 2887 2888 2889 2890 2891 2892 2893 2894 2895 2896 2897 2898 2899 2900 2901 2902 2903 2904 2905 2906 2907 2908 2909 2910 2911 2912 2913 2914 2915 2916 2917 2918 2919 2920 2921 2922 2923 2924 2925 2926 2927 2928 2929 2930 2931 2932 2933 2934 2935 2936 2937 2938 2939 2940 2941 2942 2943 2944 2945 2946 2947 2948 2949 2950 2951 2952 2953 2954 2955 2956 2957 2958 2959 2960 2961 2962 2963 2964 2965 2966 2967 2968 2969 2970 2971 2972 2973 2974 2975 2976 2977 2978 2979 2980 2981 2982 2983 2984 2985 2986 2987 2988 2989 2990 2991 2992 2993 2994 2995 2996 2997 2998 2999 3000</p>
 <p>(p. 151)</p>	
 <p>(p. 152; 217)</p>	
 <p>(p. 235)</p>	
 <p>(p. 217)</p>	
 <p>(p. 166)</p>	

(continued on next page)

Table 1 (continued)

Structure & Notes Page #	Original Image RBW Notes
 <p>(p. 167)</p>	
 <p>(p. 125)</p>	
 <p>(p. 146)</p>	
 <p>(p. 216)</p>	
 <p>(p. 262)</p>	
 <p>(p. 216)</p>	
 <p>(p. 357)</p>	
 <p>(p. 358; 361)</p>	
 <p>(p. 513)</p>	
 <p>(p. 600)</p>	




Fig. 9. Some selected structures from Fig. 8 from the RBW notes.




Fig. 10. Heterocycle reported by Woodward and co-workers in Ref. 8.




Fig. 11. Structure 20.




Fig. 12. Heterocycle reported in Ref. 8.





Fig. 13. Isoelectronic seven-membered ring heterocycles.

Networks of the 24-membered heterocycle were also considered (and beautifully rendered; see Fig. 16). The electronic structure of the infinite network of Fig. 16, $C_6S_6N_{12}$, is the last structure studied theoretically in the Whangbo, Woodward, and Hoffmann paper already mentioned.⁵

The networks envisioned by Woodward share bonds. It is tempting for the authors to speculate on a different network bonding at the C of $-N=C-N-$ moiety to give **25** (Fig. 17).

Fig. 14. 24-Membered heterocycle **24**; RBW notes, p 342.

The potential stacking scenario of the 24-membered heterocycle was also considered by Woodward (see Fig. 18). By virtue of the increased electronegativity of nitrogen in comparison to carbon and sulfur, the positive charges can be expected to reside preferentially on sulfur and carbon atoms and the negative charges on the nitrogen. Therefore, these 24-membered rings can be expected to exhibit good stacking properties, as shown in **26** and Fig. 18.

In the above hypothetical $R_6C_6N_{12}S_6$ molecule, it should be pointed out that the six sulfur atoms fall on the vertices of a regular hexagon. Again using the principles of geometry and trigonometry with the known values of the $C=N$ and $C-N$ bond lengths and the $-N-C=N-$ bond angle (120°), the distance between S_1 and S_2




Fig. 15. RBW notes, p 82, 117, 341.




Fig. 16. Networks of the 24-membered heterocycle; RBW notes, p 155, 173, 348, 351, 380.




Fig. 17. Alternate network bonding.




Fig. 18. RBW notes, p 341.

atoms can be calculated; it is near 4.6 Å. Therefore the distance between S_1 and S_4 (diagonal of the hexagon) is approximately 9.2 Å. If one places a benzene ring on the center of the 24-membered ring, the distance between S_1 and the aromatic hydrogen is computed as follows. The distance between H_1 and H_4 in the benzene ring is $2x$ the C–H bond distance (2.2 Å) plus the C_1 – C_4 distance (2.84 Å) = 5.04 Å. Since the S_1 – S_4 distance is 9.2 Å, S_1 – H_1 and S_4 – H_4 are only 2.08 Å apart. This value is considerably less than the sum of the S–H van der Waals radius (1.85+1.1=2.85 Å). Thus a benzene molecule can easily intercalate into the cavity of the 24-membered ring. A ring current can be envisaged to flow across systems, such as **27** subjected to oxidation (Fig. 19).

Hypothetical structure **15** is not known. However in 1988, almost nine years after Woodward's death, a basic seven-membered ring system viz. 1,3,5,2,4,6-trithiatiazepine **28** was synthesized by Rees (**28**, R=H Fig. 20). The seven-membered ring has been shown to undergo nitration and bromination on the single carbon atom. This may afford a route to further manipulation.²⁰

In 1985, calculations were carried out on one of the hypothetical components, viz. of trithiatetrazepine in **15**. A literature survey failed to reveal any independent attempt at the synthesis of many of the related structures depicted in **14**–**17** and a related paper describes the synthesis of some new dithiadiazenes and trithiadiazenes derived from *m*-dimercaptobenzene, as shown in Fig. 21.²²

The use of excess *m*-benzenesulfonyl chloride (**32**) led to a red polymeric material in 35% yield (Fig. 22).

The same protocol can now be used to synthesize polymer-related to **18**, using the *p*-disulfonyl chloride (Fig. 23).

Other related, hypothetical molecules conceptualized by RBW are shown in structures **38**–**41** (Fig. 24).

Woodward sketched out a possible synthetic route to such molecules shown below, using the known bis-trimethylsilylsulfurdimide **43** and a pentachloroantimonycyanide complex **42** (Fig. 25).

The resulting diamino derivative **45** can now be reacted with sulfur dichloride to give the heterocycles, **46**, **47**, etc. (Fig. 26).




Fig. 19. RBW notes, p 341, 350.




Fig. 20. Heterocycle prepared by Rees.




Fig. 23. Some reactions of molecule 34, leading to structure 37.



Fig. 21. Dithiadiazenes and trithiadiazenes derived from *m*-dimercaptobenzene (Ref. 22).

Fig. 22. Structures 29, 32, and 33.

Other interesting variations on the (SN)_x structure are those incorporating triazine and pyrimidine ring systems. These are as yet unknown, but after the death of Woodward other groups have investigated related structures independently. These are discussed next.

A class of polymers referred to as polythiocyanogens have been known for some time, and are denoted by the general formula S_Y(CN)₂, with Y being an integer.^{23,24} All of these entities polymerize very readily in solution and in the solid state to give orange and red products, which are assigned the [S_Y(CN)₂]_x formula.²⁵ As early as 1962, one assignment of the structure of [S_Y(CN)₂]_x was based on analogy. Since trimerizes to triazine, these polymers derived from polythiocyanogens were thought to be triazine networks linked by S_Y as shown in 48 (Fig. 27).




Fig. 24. Structure 38 (RBW notes, p 380); 39–41 (RBW notes, p 448).





Fig. 25. RBW notes, p 457.




Fig. 26. Structures 46 and 47.




Fig. 29. Polymer 50.







Fig. 27. Polymer 48.



51



52, R = t-Bu, Ph, etc.



53

Fig. 30. Molecules 51–53.

More recently, some thiocyanogens were made and their polymerization was studied using FTIR spectroscopy.^{25,26} An authentic sample of an oligomer **48** ($y=2$) was made by reacting trichlorotriazine and disodium disulfide (Fig. 28). This material was not identical to parathiocyanogen derived by either chemical or electrochemical oxidation of thiocyanate anion.




Fig. 28. Polymerization of 49.

A series of polythiocyanogens $[S_y(CN)_2]$ bearing 1–4 sulfur atoms per unit has been reported. These were assigned structure **50** consisting of two polyazomethine chains bridged by sulfur atoms (Fig. 29).


A related heterocycle, which can be expected to undergo ring-opening polymerization to give SN polymers is **51**. Hybrid molecules **52** and **53** can also be expected to give rise to CN/SN polymers. All of these are known but do not give rise to polymers (Fig. 30).²⁷

Woodward's instinctive feeling was that many of the idealized straight-chain structures written for $(SN)_2$ etc. may not be real led to the following proposal as to what will form an ideal conductor/superconductor. He suggests that a polymer bearing $=N-C(R)-N=S$ constrained in a ring as shown in **54** will allow for free electron flow while preventing kinks in the chain. Note his enthusiastic prediction of an organic superconductor for this structure (Fig. 31).

No specific ideas for the synthesis of such a polymer or a potential monomer are elaborated. The authors would like to propose that one needs to synthesize a diamine/diimine attached to a cyclic thiadiazine ring, which on reaction with sulfur dichloride or $SOCl_2$ should give the desired polymer (Fig. 32).

$(RCS)_x$ is a polymer isoelectronic with $(SN)_x$. It has already been seen that $(SN)_x$ is not amenable to modifications to improve its conductivity, stability etc. The one requisite in the hypothetical monomer unit is the presence of an extra unpaired electron in the pi system. The substitution of CR for N opens up the possibility of infinite modifications. In a similar vein, simple molecules isoelectronic with (SN) were considered. The simplest of these is the thioformyl radical $HC=S$ or the thioacyl radical $RC=S$. Some of the hypothetical polymers built on these group and envisioned by Woodward are shown in Fig. 33.

While these ideas may seem nebulous, one should draw attention to the work carried out by other investigators in this area. For instance a polymeric system analogous to **62** has been studied theoretically by Genin and Hoffmann (Fig. 34).²⁹



54




Fig. 31. Structure 54 (RBW notes, p 364).




Fig. 34. RBW notes, p 341.




Fig. 35. Two resonance structures for 65.

Interestingly enough, the 2,3-naphthalene analog **78** has been made and found to be stable. The difference has been ascribed to the greater contribution of a diradical structure in **77** compared to **78** (Fig. 39).³³




Fig. 39.

In addition, Woodward's unpublished notes contain ideas on TTF oligomers, organometallic derivatives, and many other beautiful structures. Some of these are displayed in the remaining figures and have no text annotation to explain their origin or context with respect to the current chemical literature. It is indeed our hope that the entire set of scanned images of the RBW notes will be made available to the chemical community as other chemists and theoreticians working in this area might be further stimulated by R. B. Woodward's remarkable imagination and depth of thought. For this remarkable organic chemist's fecund imagination thought up molecules that even now have immense interest and potential for our science Figs. 40–56.




Fig. 36. Resonance structures for 66.




Fig. 37. Reactions of 67.




Fig. 38. Cava approach to polymer 66.




Fig. 40. RBW notes, p 139.




Fig. 41. RBW notes, p 142.




Fig. 42. RBW notes, p 174.





Fig. 43. RBW notes, p 251.



$$r_{Mn} = 0.732 r_{Fe} \quad \sim (if r_{Fe} = 1.2, r_{Mn} = 0.88)$$

$$Li^+ = 0.68$$

$$Ag^{+2} = .89!$$

$$\text{volume of cube} = (2r_{Fe})^3$$

$$\text{portion taken up by Fe} = \frac{1}{8} \times 8 \left(\frac{4}{3} \pi r^3 \right)$$

$$\text{portion taken up by Mn} = \frac{4}{3} \pi (r_{Mn})^3 = \frac{4}{3} \pi (\sqrt{3}-1)^3 r_{Fe}^3$$

$$\frac{\text{total space filled}}{\text{volume of cube}} = \frac{\frac{4}{3} \pi (r_{Fe}^3 + (\sqrt{3}-1)^3 r_{Fe}^3)}{8 r_{Fe}^3} = \frac{4}{3} \pi \left(\frac{1 + (\sqrt{3}-1)^3}{8} \right) = .73$$

$$\frac{Ag Mn}{Au^{+3}} = .85$$

$$\frac{Au^{+3}}{Au Cr}$$

check volume of sphere

see hexagonal close packing

see cyanines

see Fe-Fe

see Mo-C

$$\frac{Na^+}{NaFe} = 0.97$$

$$Ca^{+2} = 0.99$$

$$\frac{CaMn}{CaFe}$$

$$Cr^{+2} = 0.89$$

$$\frac{Cr Mn}{Cr Fe}$$

$$Mg^{+1} = 0.82$$

$$\frac{Mg Fe}{Mg Mn}$$

$$Zn^{+1} = 0.88$$

$$\frac{Zn Fe}{Zn Mn}$$

$$V^{+2} = 0.88$$

$$\frac{V Mn}{V Fe}$$

275

Fig. 44. RBW notes, p 275.




Fig. 45. RBW notes, p 285.




Fig. 46. RBW notes, p 285A.

The n dimensional array has

	extra groups side chains (N_2^- or Cl^-)	ATOMS
n^3 metal atoms	—	Metal n^3
8 corners	3	$\begin{cases} Cl & 12 \\ N & 24 \end{cases}$
$12(n-2)$ non-corner edge sites	2	$\begin{cases} Cl & 12(n-2) \\ N & 12(n-2) \end{cases} \times 2$
$6(n-2)^2$ terminal edge sites	1	$\begin{cases} Cl & 3(n-2)^2 \\ N & 3(n-2)^2 \end{cases} \times 2$
$3n(n-1)^2$ center face patches	0	Cl $3n(n-1)^2$
$(n-2)^3$ M sites	0	metal only
$3n^2(n-1)$ bridges	0	N $6n^2(n-1)$

for $n=3$

$$\sum Cl = [12 + 12(n-2) + 3(n-2)^2] + 3n(n-1)^2$$

$$\sum N = 2[12 + 12(n-2) + 3(n-2)^2] + 6n^2(n-1)$$

$$\sum metal = n^3$$

for $n=3$

$$Cl = 12 + 12 + 3 + 36 = 63$$

$$N = 2(12 + 12 + 3) + 108 = 162$$

$$Cr = 27$$

N/Cr ratio is correct but there is a defect of 18 chlorine

for $n=4$

$$Cl = 12 + 24 + 12 + 108 = 156$$

$$N = 2(12 + 24 + 12) + 288 = 384$$

$$Cr = 64$$

N/Cr ratio is correct but there is a defect of 36 chlorine

for $n=5$

$$Cl = 12 + 36 + 27 + 240 = 315$$

$$N = 2(12 + 36 + 27) + 600 = 750$$

$$Cr = 125$$

N/Cr ratio is correct but there is a defect of 60 chlorine

for $n=2$ (P=6) \therefore the center faces must be being counted incorrectly.



(289)

Fig. 47. RBW notes, p 287.

$Cl^- + \begin{matrix} \text{---} \\ \text{---} \end{matrix} N_2^+ \rightarrow \begin{matrix} \text{---} \\ \text{---} \end{matrix} Cl + N_2$

$4- + 1$
 $net 3-$

$6n^2 \left[\frac{(n-1)(n-1)}{n} \right]$
 $6n \left[\frac{n(n-1) - (n-1)}{n} \right]$

$3n(n-1) [(n-1)^+]$

\therefore FINALLY

$$\sum Cl = [12 + 12(n-2) + 3(n-2)^2] + 3n(n-1)^2 + 3n(n-1)$$

$$\sum N = 2[12 + 12(n-2) + 3(n-2)^2] + 6n^2(n-1)$$

$$\sum Cr = n^3$$

$6n^2(n-1) - 6n(n-1)$ N_2 's displaced by Cl^-

Now, in fact the excess "free" ionic chlorides should be substituted at edge or face position to give an entirely neutral polycrystal, of the stoichiometry of order n

$$\sum Cl = [12 + 12(n-2) + 3(n-2)^2] + 3n^2(n-1) \leq 3n^3$$

$$\sum N = 2[12 + 12(n-2) + 3(n-2)^2] + 6n^2 \left[\frac{(n-1)^2}{n} \right]$$

$$\sum Cr = n^3$$

general metal

now as n becomes large

$$\left[\frac{(n-1)^2}{n-1} \right] = \frac{(n-1)^2}{n(n-1)} = \frac{n-1}{n}$$

n is the number of atoms along 1 edge

Lim $\frac{n-1}{n} \rightarrow 1$ and $N/Cl \rightarrow 2$

and is therefore the length of each chain, in number of metal atoms.

312

Fig. 48. RBW notes, p 312.




Fig. 49. RBW notes, p 455.




Fig. 50. RBW notes, p 533.




Fig. 51. RBW notes, p 586.




Fig. 52. RBW notes, p 638. This page clearly of lower quality than other pages, and in a different hand, is actually by Roald Hoffmann. It came out of their mutual discussions in the course of the research.




Fig. 53. RBW notes, p 641.




Fig. 54. RBW notes, p 657.




Fig. 55. RBW notes, p 676.




Fig. 56. RBW notes, p 699; the last page.

Acknowledgements

MPC and MVL wish to thank Prof. Robert M. Williams, Colorado State University for getting us involved in this challenging project. All of the authors wish to express their sincere gratitude to Mr. Eric Woodward and Ms. Crystal Woodward who made their father's notes available to us. Support from the National Science Foundation (Che-9901177 to MPC) is gratefully acknowledged. The authors are deeply indebted to Prof. Fred Wudl, of the University of California, Santa Barbara, for careful reading of the manuscript and helpful suggestions.

References and notes

- Ferraris, J.; Cowan, D. O.; Walatka, V., Jr.; Perlstein, J. H. *J. Am. Chem. Soc.* **1973**, *95*, 948.
- (a) Walatka, V. V.; Labes, M. M.; Perlstein, J. H. *Phys. Rev. Lett.* **1973**, *31*, 1139; (b) Becke-Goehring, M.; Voight, D. *Naturwissenschaften* **1953**, *40*, 482.
- Berets, D. J.; Smith, D. S. *Trans. Faraday Soc.* **1968**, *64*, 823.
- Shirakawa, H.; Louis, E. U.; MacDiarmid, A. G.; Chiang, C. K.; Heeger, A. J. *J. Chem. Soc., Chem. Commun.* **1977**, 578.
- Whangbo, M. H.; Hoffmann, R.; Woodward, R. B. *Proc. R. Soc. London A* **1979**, *366*, 23.
- Little, W. A. *Synth. Met.* **1965**, *212*; Little, W. A. *Phys. Rev.* **1964**, *134*, A1416.
- Street, G. B.; Gill, W. D.; Geiss, R. H.; Greene, R. L.; Mayerle, J. J. *J. Chem. Soc., Chem. Commun.* **1977**, 407; Gill, W. D.; Bludau, W.; Geiss, R. H.; Grant, P. M.; Greene, R. L.; Mayerle, J. J.; Street, G. B. *Phys. Rev. Lett.* **1977**, *38*, 1305; Akhtar, M.; Chiang, C. K.; Heeger, A. J.; MacDiarmid, A. G. *J. Chem. Soc., Chem. Commun.* **1977**, 846.
- Ernest, I.; Holick, W.; Rihs, G.; Schomburg, D.; Shoham, G.; Wenkert, D.; Woodward, R. B. *J. Am. Chem. Soc.* **1981**, *103*, 1540.
- (a) Boutique, J. P.; Riga, J.; Verbist, J. J.; Delhalle, J.; Fripiat, J. G.; Haddon, R. C.; Kaplan, M. L. *J. Am. Chem. Soc.* **1984**, *106*, 312; (b) Gleiter, R.; Bartetzko, R.; Cremer, D. *J. Am. Chem. Soc.* **1984**, *106*, 3437; (c) Klein, H. P.; Oakley, R. T.; Michl, J. *Inorg. Chem.* **1986**, *25*, 3194; (d) Boutique, J. P.; Lazzaroni, R.; Riga, J.; Verbist, J.; Pfister-Guillouzo, G.; Delhalle, J.; Fripiat, J. G.; Haddon, R. C.; Kaplan, M. D. *J. Chem. Soc., Perkin Trans. 2* **1987**, 9580.
- (a) Boere, R. T.; Moock, K. H.; Derrick, S.; Hoogerdijs, W.; Preuss, K.; Yip, J.; Parvez, M. *Can. J. Chem.* **1993**, *71*, 673; (b) Amin, M.; Rees, C. W. *J. Chem. Soc., Perkin Trans. 1* **1989**, 2495.
- Del, B. B.; Cordes, A. W.; Kristof, E. M.; Kristof, P.; Liblong, S. W.; Oakley, R. T. *J. Am. Chem. Soc.* **1989**, *111*, 9276.
- (a) Gimarc, B. M.; Scott, W. D. *Inorg. Chem.* **1991**, *30*, 3276; (b) Preston, K. F.; Charland, J. P.; Sutcliffe, L. H. *Can. J. Chem.* **1988**, *66*, 1299; (c) Fairhurst, S. A.; Preston, K. F.; Sutcliffe, L. H. *Phosphorus, Sulfur Silicon Relat. Elem.* **1994**, *93–94*, 105; (d) Gleiter, R.; Bartetzko, R.; Hofmann, P. Z. *Naturforsch., B: J. Chem. Sci.* **1980**, *35B*, 1166.
- Wang, L. J.; Mezey, P. G.; Zgierski, M. Z. *Chem. Phys. Lett.* **2003**, *369*, 386.
- (a) Ono, K.; Tanaka, S.; Yamashita, Y. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 1977; (b) Yamashita, Y.; Ono, K.; Tomura, M.; Tanaka, S. *Tetrahedron* **1997**, *53*, 10169; (c) Strassner, T.; Fabian, J. J. *Phys. Org. Chem.* **1997**, *10*, 33.
- Haddon, R. C.; Kaplan, M. L.; Marshall, J. H. *J. Am. Chem. Soc.* **1978**, *100*, 1235.
- (a) Boutique, J. P.; Pfister-Guillouzo, G.; Riga, J.; Verbist, J. J.; Fripiat, J. G.; Delhalle, J.; Haddon, R. C.; Kaplan, M. L. *J. Electron Spectrosc. Relat. Phenom.* **1984**, *34*, 199; (b) Folkesson, B. *J. Mol. Catal.* **1983**, *20*, 111; (c) Boutique, J. P.; Riga, J.; Verbist, J. J.; Delhalle, J.; Fripiat, J. G.; Andre, J. M.; Haddon, R. C.; Kaplan, M. L. *J. Am. Chem. Soc.* **1982**, *104*, 2691; (d) Gieren, A.; Lamm, V.; Haddon, R. C.; Kaplan, M. L. *J. Am. Chem. Soc.* **1979**, *101*, 3306; (e) Gieren, A.; Lamm, V.; Haddon, R. C.; Kaplan, M. L. *J. Am. Chem. Soc.* **1979**, *101*, 7277.
- Komin, A. P.; Street, R. N.; Carmack, M. *J. Org. Chem.* **1975**, *40*, 2749.
- (a) Bird, C. W. *Tetrahedron* **1987**, *43*, 4725; (b) Tajiri, A.; Winkler, J. Z. *Naturforsch.* **1983**, *38A*, 1263; (c) Kane, J.; Schaeffer, R. *Cryst. Struct. Commun.* **1981**, *10*, 1403; (d) Kwan, C. L.; Carmack, M.; Kochi, J. K. *J. Phys. Chem.* **1976**, *80*, 1786.
- Gimarc, B. M.; Trinajstić, J. A. *Inorg. Chim. Acta* **1985**, *102*, 105.
- Dunn, P.; Morris, J. L.; Rees, C. W. *J. Chem. Soc., Perkin Trans. 1* **1988**, 1745.
- Cramer, R. J. *Org. Chem.* **1961**, *26*, 3476.
- (a) Longridge, J. J.; Rawson, J. M. *Phosphorus, Sulfur Silicon Relat. Elem.* **1997**, *124 and 125*, 501; (b) Longridge, J. J.; Rawson, J. M. *Polyhedron* **1998**, *17*, 1871.
- Schaefer, F. C. In *The Chemistry of the Cyano Group*; Rappoport, Z., Ed.; Interscience: New York, NY, 1970; Chapter 7, p 239.
- Gmelins Handbuch der Anorganischen Chemie*; Springer: Berlin, 1978; Vol. 14/11, pp 161–237; System C14, TL D5.
- Cataldo, F.; Keheyani, Y. *Polyhedron* **2002**, *21*, 1825.
- Cataldo, F. *Polyhedron* **2000**, *19*, 681.
- Chivers, T.; Gates, D.; Li, X.; Manners, I.; Parvez, M. *Inorg. Chem.* **1999**, *38*, 70.
- Wudl, F.; Kruger, A. A.; Thomas, G. A. *Ann. N.Y. Acad. Sci.* **1978**, *313*, 79.
- Genin, H.; Hoffmann, R. *J. Am. Chem. Soc.* **1995**, *117*, 12328.
- Levinson, M.I. 'The Thionation reactions of Lawesson's reagent and Approaches to Carbon–Sulfur Analogs of (SN)_x,' Thesis, University of Pennsylvania, Philadelphia, 1984.
- Asselt, R. V.; Vanderzande, D.; Gelan, J.; Froehling, P. E.; Aagaard, O. *Synth. Met.* **2000**, *110*, 25.
- Raasch, M. S.; Huang, N.-Z.; Lakshmikantham, M. V.; Cava, M. P. *J. Org. Chem.* **1988**, *53*, 891.
- Cava, M. P.; Lakshmikantham, M. V. *Phosphorus, Sulfur Silicon Relat. Elem.* **1989**, *43*, 95.

Biographical sketch



Roald Hoffmann was born in 1937 in Złoczów, Poland. Having survived the war, he came to the U. S. in 1949, and studied chemistry at Columbia and Harvard Universities (Ph.D. 1962). Since 1965 he is at Cornell University, now as the Frank H. T. Rhodes Professor of Humane Letters, Emeritus. He has received many of the honors of his profession, including the 1981 Nobel Prize in Chemistry (shared with Kenichi Fukui). Had R.B.W. lived, he most surely would have shared in that award. Active also as a writer, Hoffmann has carved out his own land 'twixt science, poetry, and philosophy, through many essays and several books of poetry, plays, and nonfiction.



Robert M. Williams was born in New York in 1953 and received the B.A. degree in Chemistry in 1975 from Syracuse University (Ei-ichi Negishi). He obtained the Ph.D. degree in 1979 at MIT (W.H. Rastetter) and was a post-doctoral fellow at Harvard (1979–1980; R.B. Woodward/Yoshito Kishi). He joined Colorado State University in 1980 and was named a University Distinguished Professor in 2002. He has won several awards including the ACS Arthur C. Cope Scholar Award (2002) and the ACS Ernest Guenther Award in the Chemistry of Natural Products (2011). His interdisciplinary research program at the chemistry–biology interface is focused on the total synthesis of biomedically significant natural products, biosynthesis of secondary metabolites, studies on antitumor drug–DNA interactions, HDAC inhibitors, amino acids, and peptides.