

## The Green's function for the Hückel (tight binding) model

Ramis Movassagh, Gilbert Strang, Yuta Tsuji, and Roald Hoffmann

Citation: *Journal of Mathematical Physics* **58**, 033505 (2017); doi: 10.1063/1.4977080

View online: <https://doi.org/10.1063/1.4977080>

View Table of Contents: <http://aip.scitation.org/toc/jmp/58/3>

Published by the [American Institute of Physics](#)

---

### Articles you may be interested in

[Enhancing the conductivity of molecular electronic devices](#)

*The Journal of Chemical Physics* **146**, 092310 (2017); 10.1063/1.4972992

[Destructive quantum interference in electron transport: A reconciliation of the molecular orbital and the atomic orbital perspective](#)

*The Journal of Chemical Physics* **146**, 092308 (2017); 10.1063/1.4972572

[The Simpson's paradox in quantum mechanics](#)

*Journal of Mathematical Physics* **58**, 032101 (2017); 10.1063/1.4977784

[Nonequilibrium diagrammatic technique for Hubbard Green functions](#)

*The Journal of Chemical Physics* **146**, 092301 (2017); 10.1063/1.4965825


[Conformal invariance and new exact solutions of the elastostatics equations](#)

*Journal of Mathematical Physics* **58**, 031502 (2017); 10.1063/1.4977512

[Exact analytical solutions of a two-dimensional hydrogen atom in a constant magnetic field](#)

*Journal of Mathematical Physics* **58**, 042102 (2017); 10.1063/1.4979618

---



## The Green's function for the Hückel (tight binding) model

 Ramis Movassagh,<sup>1,a)</sup> Gilbert Strang,<sup>2</sup> Yuta Tsuji,<sup>3</sup> and Roald Hoffmann<sup>4</sup>
<sup>1</sup>*Department of Mathematics, IBM T.J. Watson Research Center, Yorktown Heights, New York 10598, USA*
<sup>2</sup>*Department of Mathematics, Massachusetts Institute of Technology, Cambridge Massachusetts 02139, USA*
<sup>3</sup>*Education Center for Global Leaders in Molecular Systems for Devices, Kyushu University, Nishi-ku, Fukuoka 819-0395, Japan*
<sup>4</sup>*Department of Chemistry and Chemical Biology, Cornell University, Ithaca, New York 14853, USA*

(Received 14 May 2016; accepted 9 February 2017; published online 7 March 2017)

Applications of the Hückel (tight binding) model are ubiquitous in quantum chemistry and solid state physics. The matrix representation of this model is isomorphic to an unoriented vertex adjacency matrix of a bipartite graph, which is also the Laplacian matrix plus twice the identity. In this paper, we analytically calculate the determinant and, when it exists, the inverse of this matrix in connection with the Green's function,  $\mathbf{G}$ , of the  $N \times N$  Hückel matrix. A corollary is a closed form expression for a Harmonic sum (Eq. (12)). We then extend the results to  $d$ -dimensional lattices, whose linear size is  $N$ . The existence of the inverse becomes a question of number theory. We prove a new theorem in number theory pertaining to vanishing sums of cosines and use it to prove that the inverse exists if and only if  $N + 1$  and  $d$  are odd and  $d$  is smaller than the smallest divisor of  $N + 1$ . We corroborate our results by demonstrating the entry patterns of the Green's function and discuss applications related to transport and conductivity. *Published by AIP Publishing.* [<http://dx.doi.org/10.1063/1.4977080>]

### I. PROBLEM STATEMENT: THE HÜCKEL MODEL AND ITS GREEN'S FUNCTION

The Hückel or tight binding model was originally introduced to describe electron hopping on a one-dimensional chain or ring.<sup>13</sup> It has come to serve as a ubiquitous model in solid state chemistry and physics.<sup>3,13</sup> Two typical forms of the Hückel matrix, for a linear chain of  $N$  atoms, and for a cycle of  $N$  atoms, are given in Eq. (3). The resulting banded matrix is isomorphic to the vertex adjacency matrix of a graph.<sup>12</sup>

The diagonal entries of the Hückel Hamiltonian matrix are defined by the Coulomb integral,  $\alpha = \langle \chi_i | \hat{H} | \chi_i \rangle$ , where  $\chi_i$  is the basis function of  $2p_z$  AO of the  $i$ th carbon atom. The magnitude of  $\alpha$  can be approximated by the ionization potential of a carbon atom. It is reasonable to assume that all carbon atoms have the same ionization potential, resulting in an approximation that all diagonal elements have the same value, which is denoted by just  $\alpha$ .

An off-diagonal element of the Hückel Hamiltonian matrix is typically called the resonance integral,  $\beta_{i,j}$ , defined as  $\beta_{i,j} = \langle \chi_i | \hat{H} | \chi_j \rangle$ . This is a measure of the interaction between the  $i$ th and  $j$ th carbon atoms. Usually  $\beta_{i,j}$  is neglected if there is no bond between the  $i$ th and  $j$ th carbon atoms. It is reasonably assumed that all the carbon-carbon bonds have the same strength, resulting in an approximation that all non-zero off-diagonal elements have the same value, which is denoted by just  $\beta$ .

If there are no heteroatoms in the system, all Coulomb integrals can be considered identical. However, if there is a heteroatom, we need to modify the Coulomb integral properly. When one calculates a linear  $\pi$ -conjugated chain, namely, a polyene, all resonance integrals cannot be considered

<sup>a)</sup>Electronic address: [ramis.mov@gmail.com](mailto:ramis.mov@gmail.com)



where  $C_{rk}$  is the coefficient of the  $r$ th atomic orbital (AO) in the  $k$ th molecular orbital (MO) in an orthogonal basis,  $\epsilon_k$  is the  $k$ th MO energy, and  $\eta$  is an infinitesimal positive number to assure analyticity. The Fermi energy is equal to the Coulomb integral of the Hückel model, and for convenience we set this energy to zero (see Subsection I B).

For the DC conductivity, we want to evaluate the Green's function at the Fermi energy; therefore  $E = 0$  in Eq. (4). By the Sokhotski–Plemelj theorem,  $\lim_{\eta \rightarrow 0^+} \frac{1}{x \pm i\eta} = P(\frac{1}{x}) \mp i\pi\delta(x)$ , the real part of the Green's function (Eq. (5)) is

$$G^{(0)}(r, s; E = 0) = - \sum_k \frac{C_{rk} C_{sk}^*}{\epsilon_k}. \quad (6)$$

Therefore, the Green's function in the basis described above (Eqs. (1) and (2)) for a finite open linear chain has entries that are

$$G^{(0)}(r, s; E = 0) = - \frac{1}{N+1} \sum_{k=1}^N \frac{\sin(rk\omega) \sin(sk\omega)}{\cos(k\omega)} \quad \text{in units of } \beta^{-1}, \quad (7)$$

$$\omega \equiv \frac{\pi}{N+1}.$$

Below, for simplicity, we denote  $G(r, s) \equiv G^{(0)}(r, s; E = 0)$ .

*Remark 1.*  $\mathbf{G}^{(0)}$  is simply  $-H_1^{-1}$  in the basis given by Eqs. (1) and (2). Generally, with energy set-point  $E = 0$ , the Green's function is minus the inverse of the Hamiltonian. We compute the inverses of  $H_1$  and  $H_1^c$  in several ways—from general formulas for tridiagonal matrices, directly from simple equations for the first column of the inverses, and from factoring the matrix symbol  $e^{i\theta} + e^{-i\theta}$ .

Our goal is to prove the conditions under which the inverse of various forms of the Hückel model exists for different  $N$  and in  $d$ -spatial dimensions. When it exists, we analytically derive closed-form formulas for the Green's function  $\mathbf{G}^{(0)}$ .

## B. Level set of the Fermi energy

The position of the actual Fermi levels in a calculation of molecular transmission may vary. It has proven to be a good approximation to set it equal to the Coulomb integral of the Hückel method for most rings and chains (the limitations of this assumption will be mentioned later).

There is a good reason why we assume  $E_F = \alpha$ . The energy level of the  $2p$  atomic orbital (AO) of carbon ( $-11.4$  eV, the same energy level as the Coulomb integral), is almost the same as that of the  $6s$  AO of the widely used Au electrode ( $-10.9$  eV).<sup>2</sup> Note that the electronic configuration of Au is  $5d^{10}6s^1$ . This approximation works well as long as significant charge transfer between the molecule and the electrode surface does not occur.<sup>30</sup>

The assumption that the Fermi level is equal to the Coulomb integral of the Hückel method is probably valid for even-membered chains and rings with  $4n + 2$  atoms.

## II. DETERMINANTS AND ANALYTICAL EXPRESSIONS FOR $H_1^{-1}$ AND $(H_1^c)^{-1}$

### A. Open chain, $H_1$

*Lemma 1.*  $H_1$  is only invertible when  $N$  is even, in which case  $\det(H_1) = (-1)^{N/2}$ .

*Proof.* Generally for any  $N$ ,

$$\det(H_N) = -\det(H_{N-2}) = \dots = (-1)^{\frac{N-2}{2}} \det(H_2) = (-1)^{\frac{N-2}{2}} \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} = (-1)^{N/2} \quad N \text{ even,}$$

$$\det(H_N) = -\det(H_{N-2}) = \dots = \det(H_1) = |0| = 0 \quad N \text{ odd,}$$

where we denoted  $H_1$  of size  $N \times N$ , simply by  $H_N$  and the determinant of a matrix by  $|\cdot|$ . In Eq. (7),  $\cos(k\omega)$  can take a zero value if  $N$  is odd, whereby  $G(r, s)$  is not defined.  $\square$

*Proposition 1.* When  $N$  is even, the entries of the Green's function  $G(r, s) \equiv -H_1^{-1}(r, s)$  are

$$G(r, s) = \begin{cases} (-1)^{\frac{r+s-1}{2}} & r < s : r \text{ odd and } s \text{ even} \\ (-1)^{\frac{r+s-1}{2}} & r > s : r \text{ even and } s \text{ odd}, \\ 0 & \text{otherwise} \end{cases}, \tag{8}$$

$$G = -H_1^{-1} = \begin{bmatrix} 0 & -1 & 0 & +1 & 0 & -1 & \dots \\ -1 & 0 & 0 & 0 & 0 & 0 & \\ 0 & 0 & 0 & -1 & 0 & +1 & \\ +1 & 0 & -1 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & -1 & \\ -1 & 0 & +1 & 0 & -1 & 0 & \dots \\ \vdots & & \vdots & & & & \ddots \end{bmatrix}. \tag{9}$$

*Proof.* Suppose we have a general tridiagonal matrix

$$A = \begin{bmatrix} b_1 & c_1 & & & & \\ a_1 & b_2 & c_2 & & & \\ & a_2 & b_3 & \ddots & & \\ & & \ddots & \ddots & c_{N-1} & \\ & & & a_{N-1} & b_N & \end{bmatrix},$$

then Usmani's formula<sup>27</sup> for the  $r, s$  entry of  $A^{-1}$  is

$$\alpha(r, s) = \begin{cases} (-1)^{r+s} c_r c_{r+1} \dots c_{s-1} \theta_{r-1} \phi_{s+1} / \theta_N & r < s \\ \theta_{r-1} \phi_{r+1} / \theta_N & r = s, \\ (-1)^{r+s} a_{s+1} a_{s+2} \dots a_r \theta_{s-1} \phi_{r+1} / \theta_N & r > s \end{cases}, \tag{10}$$

where  $\theta_r$  and  $\phi_s$  satisfy second order recursion relations

$$\begin{aligned} \theta_r &= b_r \theta_{r-1} - a_r c_{r-1} \theta_{r-2}, \quad r = 1, 2, \dots, N-1, N, \\ \phi_s &= b_s \phi_{s+1} - c_s a_{s+1} \phi_{s+2}, \quad s = N, N-1, \dots, 2, 1, \end{aligned}$$

with the initial conditions  $\theta_{-1} = 0, \theta_0 = 1, \phi_{N+1} = 1$ , and  $\phi_{N+2} = 0$ .

We are interested in the special case where  $b_i = 0, c_i = a_i = 1$  for all  $i$ . The recursion relations are now given by

$$\begin{aligned} \theta_r &= -\theta_{r-2}, \quad r = 2, 3, \dots, N, \\ \phi_s &= -\phi_{s+2}, \quad s = N, N-1, \dots, 2, 1. \end{aligned}$$

The solutions, after imposing the initial conditions, are

$$\theta_r = \frac{i^r}{2} [1 + (-1)^r], \quad \phi_s = \frac{i^{-(N+1)+s}}{2} [1 - (-1)^s].$$

Substituting these in Eq. (10) and multiplying by  $-1$ , we obtain  $\mathbf{G} = -H_1^{-1}$ ,

$$G(r, s) = \begin{cases} \frac{i^{3(r+s-1)}}{4} [1 + (-1)^{r-1}] [1 - (-1)^{s+1}] & r < s \\ \frac{i^{2(r-1)}}{4} [1 + (-1)^{r-1}] [1 - (-1)^{r+1}] & r = s, \\ \frac{i^{3(r+s-1)}}{4} [1 + (-1)^{s-1}] [1 - (-1)^{r+1}] & r > s \end{cases}, \tag{11}$$

$\square$

By symmetry we may focus on  $r \geq s$ . In Eq. (11) the only nonzero elements, for  $r \geq s$ , correspond to  $r$  even and  $s$  odd, in which case  $G(r, s) = i^{3(r+s-1)} = (-1)^{\frac{r+s-1}{2}}$ .



In particular when  $n = 4j$ , the matrix has zero eigenvalues and hence non-invertible. The following lemma sharpens this notion.

*Lemma 2.* The determinant of  $H_1^c$  is given by

$$\det(H_1^c) = \begin{cases} -1 & N = 2 \\ 2 & N = 2k + 1 \\ 0 & N = 4k \\ -4 & N = 4k + 2 \end{cases}, \quad \text{for } k \in \mathbb{N}.$$

*Proof.* When  $N = 2$ , trivially  $\begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} = -1$ . When  $N$  is odd, we express

$$H_1^c = \begin{bmatrix} H_{N-1} & B \\ C & D \end{bmatrix}$$

where  $C = [1 \ 0 \ \dots \ 0 \ 1]$ ,  $B=C^T$ ,  $D = 0$  and  $H_{N-1}$  is an  $(N - 1) \times (N - 1)$  version of  $H_1$  (open chain) as defined above, which is invertible. With this decomposition, the structure of  $H_1^{-1}$  derived above, and the well-known fact about the determinant of block matrices we arrive at

$$\det(H_1^c) = -\det(H_{N-1}) \det(CH_{N-1}^{-1}B) = \det(CH_{N-1}^{-1}B) = 2.$$

When  $N$  is a multiple of 4, one can easily check that the vectors  $\mathbf{v}_1 \equiv [0, -1, 0, 1]^T$  and  $\mathbf{v}_2 \equiv [1, 0, -1, 0]^T$  generate the kernel of  $H_1^c$ . Namely, if  $H_1^c$  is a  $4k \times 4k$  matrix, then the  $k$ -fold concatenations  $[\mathbf{v}_1 \mathbf{v}_1 \dots \mathbf{v}_1]^T$  and  $[\mathbf{v}_2 \mathbf{v}_2 \dots \mathbf{v}_2]^T$  are in the  $\ker(H_1^c)$ . Moreover, since excluding the last two rows and columns of  $H_1^c$  gives  $H_{4k-2}$ , which is invertible, we conclude that the two vectors are a basis for the kernel of  $H_1^c$ .

Lastly, if  $N$  is even yet not a multiple of 4, we write  $H_1^c = \begin{bmatrix} H_{N-2} & B \\ B^T & D \end{bmatrix}$ , where  $B^T = \begin{bmatrix} 0 \dots 0 & 1 \\ 1 & 0 \dots 0 \end{bmatrix}$  and  $D = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ . Here  $H_{N-2}$  has a size that is a multiple of 4. Using the techniques above, we obtain

$$\begin{aligned} \det(H_1^c) &= \det(H_{N-2}) \det(D - B^T H_{N-2}^{-1} B) = \det(D - B^T H_{N-2}^{-1} B) \\ &= \det\left(\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} - \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}\right) = -4. \end{aligned}$$

□

*Definition 2.* A Toeplitz matrix is a matrix that is constant along diagonals. A circulant matrix is Toeplitz, and each column is a cyclic shift of the previous column.<sup>11,23</sup> Thus the lower triangular part of a circulant determines the upper triangular part,

$$\begin{aligned} \text{Toeplitz } A &= \begin{bmatrix} x_0 & x_{-1} & x_{-2} \\ x_1 & x_0 & x_{-1} \\ x_2 & x_1 & x_0 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} x_0 & x_{-1} & \dots & x_{-(N-1)} \\ x_1 & x_0 & & \vdots \\ \vdots & & \ddots & x_{-1} \\ x_{N-1} & \dots & x_1 & x_0 \end{bmatrix}, \\ \text{Circulant } A &= \begin{bmatrix} x_0 & x_2 & x_1 \\ x_1 & x_0 & x_2 \\ x_2 & x_1 & x_0 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} x_0 & x_{N-1} & \dots & x_1 \\ x_1 & x_0 & & x_2 \\ \vdots & & \ddots & \vdots \\ x_{N-1} & x_{N-2} & \dots & x_0 \end{bmatrix}. \end{aligned}$$

The inverse of a circulant matrix is circulant. The inverse of a Toeplitz matrix is not in general Toeplitz.





$$\begin{aligned}
 N = 4k + 1 \quad x_{N-2} = x_{4k-1} &= -\frac{1}{2} \text{ by Eq. (14) and then } x_0 = \frac{1}{2}, \\
 N = 4K + 2 \quad x_{N-2} = x_{4k} = x_0 &\text{ by Eq. (15) and then } x_0 = -x_0 = 0, \\
 N = 4k + 3 \quad x_{N-2} = x_{4k+1} &= +\frac{1}{2} \text{ by Eq. (14) and then } x_0 = -\frac{1}{2}.
 \end{aligned}$$

The alternating signs for  $x_0, x_2, x_4, \dots$  complete the inverse circulant matrix  $(H_1^c)^{-1}$ . The Green's matrix is defined as  $G_1^c \equiv -(H_1^c)^{-1}$  and is shown in Fig. 1 □

*Remark 3.* The same direct approach produces  $H_1^{-1}$  in the non-circulant case. The  $(1, n)$  and  $(n, 1)$  entries of  $H_1^c$  are now set to zero. Then the first equation in Eq. (14) is simply  $x_1 = 1$ . The other equations in Eq. (14) give alternating signs for  $x_3, x_5, \dots$

Similarly, the last equation in Eq. (15) is now  $x_{N-2} = 0$ . The first column  $(x_0, x_1, \dots)$  of  $H_1^{-1}$  is seen to be  $(0, 1, 0, -1, \text{repeat})$ . The last column of  $H_1^{-1}$  has these components in the reverse order. Then by symmetry we also know the first and last rows of  $H_1^{-1}$ .

Because  $H_1$  is tridiagonal, these two columns and two rows completely determine the rest of  $H_1^{-1}$ . On and above the main diagonal, all sub-matrices of  $H_1^{-1}$  have rank 1 (If  $H_1$  is tridiagonal and invertible then  $H_1^{-1}$  is a "semi-separable" matrix.<sup>28</sup>). It is easy to see that starting from the first and last rows and columns of  $G$  in Proposition 2, all other entries of  $G$  follow directly from the rank 1 requirement.

*Proof.* (alternative to Proposition 3) We now establish the inverse of  $H_1^c$  using a technique that is general to circulant matrices based on the factorization of the symbol.  $H_1^c = S + S^{-1} = S + S^T$ , where  $S$  is the  $N \times N$  cyclic shift matrix:  $S^N = I$ . Since  $S + S^{-1} = (\mathbb{I} - iS)(\mathbb{I} + iS)S^{-1}$ , we have

$$\begin{aligned}
 (S + S^{-1})^{-1} &= S(\mathbb{I} + iS)^{-1}(\mathbb{I} - iS)^{-1} \\
 &= S \frac{(\mathbb{I} + (-i)S + \dots + (-i)^{N-1}S^{N-1})}{1 - (-i)^N} \frac{(\mathbb{I} + iS + \dots + i^{N-1}S^{N-1})}{1 - i^N}. \tag{16}
 \end{aligned}$$

The denominator is  $(1 - i^N)(1 - (-i)^N) = \begin{cases} 0 & N \equiv 4k \\ 2 & N \equiv 4k + 1 \\ 4 & N \equiv 4k + 2 \\ 2 & N \equiv 4k + 3 \end{cases}$ . This confirms that  $H_1^c$  is singular for

$N = 4k$ .

The coefficient of  $S^N$  in the product given by Eq. (16) is the numerator,

$$i^{N-1} + (-i)i^{N-2} + (-i)^2i^{N-3} + \dots + (-i)^{N-2}i + (-i)^{N-1} = \frac{i^N - (-i)^N}{i - (-i)} = \begin{cases} +1 & N \equiv 1 \pmod{4} \\ 0 & N \equiv 2 \pmod{4} \\ -1 & N \equiv 3 \pmod{4} \end{cases}.$$

When we divide by the denominators 2, 4, 2 we find the main diagonal of  $(H_1^c)^{-1}$  as the coefficients of  $S^N = \mathbb{I}$  in  $(S + S^{-1})^{-1}$ :  $\frac{1}{2}, 0, -\frac{1}{2}$  for  $N: 4k + 1, 4k + 2, 4k + 3$ , respectively.

Now we find the coefficient of  $S = S^{N+1}$  in Eq. (16). The numerator is  $1 + i^{N-1}(-i) + i^{N-2}(-i)^2 + \dots + i(-i)^{N-1} = 1 + (i)(-i)[i^{N-2} + i^{N-1}(-i) + \dots + (-i)^{N-2}]$ . Simplifying the numerator we find  $1 + \frac{i^{N-1} - (-i)^{N-1}}{i - (-i)} = 1, 2, -1$  for  $N: 4k + 1, 4k + 2, 4k + 3$ , respectively.

Dividing by 2, 4, 2 in the denominator, we find  $\frac{1}{2}$  on the diagonals  $\pm 1$  of  $(S + S^{-1})^{-1}$ .

Finally, notice that diagonals 2, 3, 4, 5, ... of  $(S + S^{-1})^{-1}$  will have opposite sign to diagonals 0, 1, 2, 3, ... The multiplication in the numerator of Eq. (16) gives a cyclic convolution

$$(1, i, i^2, \dots, i^{N-1}) \star (1, -i, (-i)^2, \dots, (-i)^{N-1})$$

for the coefficients of  $S, S^2, \dots$ . Because  $i^2 = -1$ , the coefficient of  $S^{k+2}$  in the numerator of Eq. (16) is the negative of the coefficient of  $S^k$ . The denominators are still 2, 4, 2 for  $N \equiv 1, 2, 3$ . So the pattern

in  $(S + S^{-1})^{-1} = -\mathbf{G}_1^c$  (starting with the main diagonal) is

$$\begin{aligned} N \equiv 4k + 1 & \text{ diagonals } \frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{2} \text{ repeated,} \\ N \equiv 4k + 2 & \text{ diagonals } 0, \frac{1}{2}, 0, -\frac{1}{2} \text{ repeated,} \\ N \equiv 4k + 3 & \text{ diagonals } -\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2} \text{ repeated.} \end{aligned}$$

This completes the alternative proof. □

Analogous to Proposition 2, the proposition below gives the form of the cyclic Hamiltonian with the special case of alternating bond strengths and its corresponding Green’s function.

*Definition 3.* The cyclic bond alternating Hamiltonian is defined by ( $N$  even)

$$H_{alt}^c = \begin{bmatrix} 0 & \beta & & & \alpha \\ \beta & 0 & \alpha & & \\ & \alpha & 0 & \beta & \\ & & \beta & 0 & \ddots \\ & & & \ddots & \ddots & \alpha \\ \alpha & & & & \alpha & 0 & \beta \\ & & & & & \beta & 0 \end{bmatrix}.$$

Comment: As before, this model is not circulant nor has it the Toeplitz structure.

*Proposition 4.* The entries of the Green’s function  $-(H_{alt}^c)^{-1}$  are given by

$$G(r, s) = -\frac{1}{4} \begin{cases} \frac{(-\alpha/\beta)^{\frac{r-s-1}{2}}}{\beta \left[1 - \left(-\frac{\alpha}{\beta}\right)^{N/2}\right]} [1 + (-1)^r] [1 - (-1)^s] + \frac{(-\beta/\alpha)^{\frac{r-s-1}{2}}}{\alpha \left[1 - \left(-\frac{\beta}{\alpha}\right)^{N/2}\right]} [1 - (-1)^r] [1 + (-1)^s] & r > s \\ \frac{(-\alpha/\beta)^{\frac{N+r-s-1}{2}}}{\beta \left[1 - \left(-\frac{\alpha}{\beta}\right)^{N/2}\right]} [1 + (-1)^r] [1 - (-1)^s] + \frac{(-\beta/\alpha)^{\frac{N+r-s-1}{2}}}{\alpha \left[1 - \left(-\frac{\beta}{\alpha}\right)^{N/2}\right]} [1 - (-1)^r] [1 + (-1)^s] & r \leq s \end{cases} \quad (17)$$

*Proof.* We obtain the inverse by solving for  $\mathbf{y}$  in  $H_{alt}^c \mathbf{x} = \mathbf{y}$ ; that is, we think of  $\mathbf{y}$  as given and we solve for  $\mathbf{x}$ . This will give us  $\mathbf{x} = (H_{alt}^c)^{-1} \mathbf{y}$ . First we solve the even rows in terms of the last row  $x_N$ , which itself can be solved from  $x_N = \sum_{i=1}^N [(H_{alt}^c)^{-1}]_{N,i} y_i$  to give

$$\begin{aligned} x_{2k} &= \frac{1}{\beta} \left\{ \sum_{m=0}^{k-1} \left(-\frac{\alpha}{\beta}\right)^m y_{2k-2m-1} \right\} + \left(-\frac{\alpha}{\beta}\right)^k x_N, \\ x_N &= \beta^{-1} \left[1 - (-\alpha/\beta)^{N/2}\right]^{-1} \sum_{m=0}^{\frac{N}{2}-1} \left(-\frac{\alpha}{\beta}\right)^m y_{N-2m-1}. \end{aligned}$$

Similarly the odd rows are obtained in terms of  $x_1$ , which itself can be solved  $x_1 = \sum_{i=1}^N [(H_{alt}^c)^{-1}]_{1,i} y_i$  to give

$$\begin{aligned} x_{2k+1} &= \frac{1}{\alpha} \left\{ \sum_{m=0}^{k-1} \left(-\frac{\beta}{\alpha}\right)^m y_{2k-2m} \right\} + \left(-\frac{\beta}{\alpha}\right)^k x_1, \\ x_1 &= \alpha^{-1} \left[1 - (-\beta/\alpha)^{N/2}\right]^{-1} \left\{ y_N - \frac{\beta}{\alpha} \sum_{m=0}^{\frac{N}{2}-1} \left(-\frac{\beta}{\alpha}\right)^m y_{N-2m-2} \right\}. \end{aligned}$$

Combining these equations to solve for the even and odd rows separately and multiplying by an overall minus sign, we arrive at  $\mathbf{G} = -(H_{alt}^c)^{-1}$  given by Eq. (17). □

Comment: In the special case that  $N = 2k(2k - 1)$ ,  $G_1^c$  in Proposition 3 can be obtained from Eq. (17) by substituting  $\alpha = \beta = 1$ . Note that in this limit, it is necessary that  $N \neq 4k$  for the denominator not to vanish in agreement with Lemma 2.

We now pose a more general (and difficult) question. When does the inverse exist in spatial dimension  $d$  and if it does, how can it be computed? In Sec. III we use mathematical techniques borrowed from quantum information theory and number theory to address some of these problems.

### III. HIGHER DIMENSIONAL GREEN'S FUNCTION

The Green's function we derived is the negative of the inverse of the Hückel (tight binding) Hamiltonian, whose  $N \times N$  matrix representation in Dirac notation<sup>7</sup> is

$$H_1 = \sum_{k=1}^N \{ |k\rangle\langle k+1| + |k+1\rangle\langle k| \} \quad , \quad (18)$$

where in units of  $\beta$  the coupling can be taken to be one.

To explore the  $d$ -dimensional analog  $H_d$ , we use tensor products of matrices. Recall that the tensor product of an  $m \times n$  matrix  $A$  and an  $p \times q$  matrix  $B$  is the  $mp \times nq$  matrix defined by

$$A \otimes B = \begin{bmatrix} a_{11}B & a_{12}B & \cdots & a_{1n}B \\ a_{21}B & a_{22}B & \cdots & a_{2n}B \\ \vdots & \vdots & & \vdots \\ a_{m1}B & a_{m2}B & \cdots & a_{mn}B \end{bmatrix} .$$

The Hamiltonian,  $H_d$ , on a square lattice in  $d$ -spatial dimensions (square lattice in  $d = 2$ , cubic in  $d = 3$ , etc.), with the linear size  $N$  can succinctly be expressed as

$$H_d = \sum_{i=1}^d \mathbb{I}_{N^{i-1}} \otimes H_1 \otimes \mathbb{I}_{N^{d-i}} \quad , \quad (19)$$

where  $H_1$  is given by Eq. (18), and the size of every identity matrix is indicated by its subscript. In dimensions 2 and 3, the Hamiltonians come from  $H_1$  and  $\mathbb{I} = \mathbb{I}_N$ ,

$$H_2 = (H_1 \otimes \mathbb{I}) + (\mathbb{I} \otimes H_1) \quad , \quad (20)$$

$$H_3 = (H_1 \otimes \mathbb{I} \otimes \mathbb{I}) + (\mathbb{I} \otimes H_1 \otimes \mathbb{I}) + (\mathbb{I} \otimes \mathbb{I} \otimes H_1) \quad . \quad (21)$$

Comment: The techniques apply more generally where the lattice can be constructed from  $d$  independent linear subsets.

Comment: When  $H_1$  is a Toeplitz or a circulant matrix, the corresponding  $H_d$  is generally not a Toeplitz or a circulant matrix,<sup>18</sup> but they will be block Toeplitz or block circulant, respectively.

The eigenvalue decomposition of  $H_1 = Q\Lambda Q^T$ , where  $\Lambda$  is the  $N \times N$  diagonal matrix of eigenvalues whose  $k$ th entry is  $2 \cos k\omega$  and  $Q$  is the matrix of eigenvectors with  $r$ th column given by Eq. (2). Since  $\cos k\omega \neq 0$  for all  $1 \leq k \leq N$ ,  $\Lambda$  is a diagonal matrix with no zero entries on the diagonal and  $H_1$  is invertible, i.e., has a Green's function, as expected from our calculations.

The associated Green's function matrix in  $d$  dimensions is defined by  $\mathbf{G}_d = -H_d^{-1}$ . Obtaining an analytical expression for the inverse in higher dimensions, at first, might seem difficult because it involves sums of matrices. In  $d = 2$  the size of the lattice is  $N \times N$  and in  $d = 3$  the size is  $N \times N \times N$ .

After the eigenvalue decomposition, the Hamiltonians in higher dimensions (e.g., Eqs. (20) and (21)) read

$$H_d = Q^{\otimes d} \left\{ \sum_{i=1}^d \mathbb{I}_{N^{i-1}} \otimes \Lambda \otimes \mathbb{I}_{N^{d-i}} \right\} (Q^T)^{\otimes d} \quad (22)$$

$$\equiv Q^{\otimes d} \{ \Lambda_d \} (Q^T)^{\otimes d} \quad , \quad (23)$$

where the matrix of eigenvectors denoted by  $Q^{\otimes d} \equiv Q \otimes \dots \otimes Q$  is a  $d$ -fold tensor product and the diagonal matrix of eigenvalues is  $\Lambda_d = \sum_{i=1}^d \mathbb{I}_{N^{i-1}} \otimes \Lambda \otimes \mathbb{I}_{N^{d-i}}$ . For example,

$$H_2 = (Q\Lambda Q^T \otimes \mathbb{I}) + (\mathbb{I} \otimes Q\Lambda Q^T) \tag{24}$$

$$= (Q \otimes Q) [(\Lambda \otimes \mathbb{I}) + (\mathbb{I} \otimes \Lambda)] (Q^T \otimes Q^T),$$

$$H_3 = (Q\Lambda Q^T \otimes \mathbb{I} \otimes \mathbb{I}) + (\mathbb{I} \otimes Q\Lambda Q^T \otimes \mathbb{I}) + (\mathbb{I} \otimes \mathbb{I} \otimes Q\Lambda Q^T) \tag{25}$$

$$= (Q \otimes Q \otimes Q) [(\Lambda \otimes \mathbb{I} \otimes \mathbb{I}) + (\mathbb{I} \otimes \Lambda \otimes \mathbb{I}) + (\mathbb{I} \otimes \mathbb{I} \otimes \Lambda)] (Q^T \otimes Q^T \otimes Q^T).$$

This change of basis allows us to diagonalize the Hamiltonians in any dimension, for example,

$$\Lambda_2 = (\Lambda \otimes \mathbb{I}) + (\mathbb{I} \otimes \Lambda), \tag{26}$$

$$\Lambda_3 = (\Lambda \otimes \mathbb{I} \otimes \mathbb{I}) + (\mathbb{I} \otimes \Lambda \otimes \mathbb{I}) + (\mathbb{I} \otimes \mathbb{I} \otimes \Lambda). \tag{27}$$

Below we investigate the conditions under which the Green’s function exists. For now suppose that it does. Its algebraic representation in  $d$  dimensions (compare with Eqs. (22) and (23)) is

$$\mathbf{G}_d = -Q^{\otimes d} \left\{ \sum_{i=1}^d \mathbb{I}_{N^{i-1}} \otimes \Lambda \otimes \mathbb{I}_{N^{d-i}} \right\}^{-1} (Q^T)^{\otimes d}. \tag{28}$$

It is clear that if  $\mathbf{G}_d$  were to exist no eigenvalue can be zero. Namely, diagonal entries being all the possible sums should satisfy  $2 \sum_{i=1}^d \cos \left( \frac{k_i \pi}{N+1} \right) \neq 0$  for any choice of  $1 \leq k_i \leq N$ . Then, the corresponding eigenvalues of  $\mathbf{G}_d$  are  $-1 / \left\{ 2 \sum_{i=1}^d \cos \left( \frac{k_i \pi}{N+1} \right) \right\}$ .

As an illustration let us take  $d = 2$ . Then the energies are the diagonal entries of  $\Lambda_2$  given by the sum

$$\Lambda_2 = 2 \begin{bmatrix} (\cos \omega) \mathbb{I} & & & \\ & (\cos 2\omega) \mathbb{I} & & \\ & & \ddots & \\ & & & (\cos N\omega) \mathbb{I} \end{bmatrix} + \begin{bmatrix} \Lambda & & & \\ & \Lambda & & \\ & & \ddots & \\ & & & \Lambda \end{bmatrix},$$

which is a matrix of size  $N^2 \times N^2$ ;  $\Lambda$  and  $(\cos k\omega) \mathbb{I}$  are  $(N \times N)$ . Since  $\cos k\omega = -\cos [(N + 1 - k)\omega]$ , each block of the sum is  $2(\cos k\omega) \mathbb{I} + \Lambda$  for some  $1 \leq k \leq N$  whose  $(N + 1 - k)$ th entry is zero. Therefore, the diagonal  $N^2 \times N^2$  matrix  $\Lambda_2$  has exactly  $N$  zeros on its diagonal, one in each of the  $N$  blocks, and hence noninvertible.

**IV. GREEN’S FUNCTION AND NUMBER THEORY**

The existence of the Green’s function,  $\mathbf{G}_d$ , in higher dimensions requires that  $H_d$  has non-zero eigenvalues, i.e.,  $\sum_{i=1}^d \cos \left( \frac{k_i \pi}{N+1} \right) \neq 0$  for any choice of  $1 \leq k_i \leq N$ .

*Lemma 3.*  $H_d^{-1}$  does not exist in even spatial dimensions.

*Proof.* Since  $\cos k\omega = -\cos [(N + 1 - k)\omega]$  for any  $1 \leq k \leq N$ , we can always pair up the cosines such that each pair sums to zero implying that there is a zero eigenvalue. □

Therefore, below we take  $d$  and  $N + 1$  to be odd (as  $N$  odd is already non-invertible in one dimension).

We need to prove the general conditions under which  $H_d$  is invertible, which is a problem in number theory. Recently there has been quite a bit of interest in a closely related question, which is under what conditions do sums of roots of unity vanish? Besides sheer theoretical interest, this problem is related to many mathematical structures. For example, Poonen and Rubinstein relate this problem to the number of interior intersection points made by the diagonals of a regular  $n$ -gon.<sup>19</sup>

Let us denote  $n \equiv N + 1$ . Suppose one asks for what natural numbers  $d$  do there exist  $n$ th roots of unity  $\alpha_1, \dots, \alpha_d \in \mathbb{C}$  such that  $\alpha_1 + \dots + \alpha_d = 0$ ? Such an equation is said to be a vanishing sum

of  $n$ th roots of unity of weight  $d$ . Let  $n$  have the prime factorization  $p_1^{a_1} \cdots p_r^{a_r}$  ( $a_i > 0$ ), then we can define  $W(n)$  to be the set of weights  $d$  for which there exists a vanishing sum  $\alpha_1 + \cdots + \alpha_d = 0$ ; if the sum does not vanish, then  $W(n)$  is simply the empty set.

Before delving into the proof we introduce some notation and terminology presented in Ref. 14. Let  $\langle G \rangle$  be a cyclic group of order  $n$  and let  $\zeta$  be a (fixed) primitive  $n$ th root of unity. There exists a natural ring homomorphism  $\varphi$  from the integral group  $\mathbb{Z}G$  to the ring of cyclotomic integers  $\mathbb{Z}[\zeta]$ , given by the equation  $\varphi(z) = \zeta$ , i.e., the map  $\varphi : \mathbb{Z}G \rightarrow \mathbb{Z}[\zeta]$ . An element of  $\mathbb{Z}G$ , say  $x = \sum_{g \in G} x_g g$ , lies in the kernel  $\ker(\varphi)$  if and only if  $\sum_{g \in G} x_g \varphi(g) = 0$  in  $\mathbb{Z}[\zeta]$ . Therefore, the elements of the ideal  $\ker(\varphi)$  correspond precisely to all  $\mathbb{Z}$ -linear relations among the  $n$ th roots of unity. For vanishing sums of  $n$ th roots of unity, we have to look at elements  $x = \sum_g x_g g \in \ker(\varphi)$  with  $x_g \geq 0$ ; the number of non-zero coefficients  $x_g$  is denoted by  $\epsilon_0(x)$ . In other words one looks at  $\mathbb{N}G \cap \ker(\varphi)$ , where  $\mathbb{N}G$  denotes the group semi-ring of  $G$  over  $\mathbb{N}$ .

A vanishing sum  $\alpha_1 + \cdots + \alpha_d = 0$  is called minimal if no proper sub-sum is zero. Clearly, one can always multiply a vanishing sum by a root of unity to get another vanishing sum; we say that the latter is similar to the former, i.e., one can be obtained from the other by a rotation. For any natural number  $n$ ,  $\zeta_n$  denotes a primitive  $n$ th root of unity in  $\mathbb{C}$ .

In terms of roots of unity, a vanishing sum from the basic relations of the form

$$1 + \zeta_{p_i} + \zeta_{p_i}^2 + \cdots + \zeta_{p_i}^{p_i-1} = 0 \quad 1 \leq i \leq r \tag{29}$$

is called a symmetric minimal elements in  $\mathbb{N}G \cap \ker(\varphi)$ . In general, there are vanishing minimal sums which are not similar to those in Eq. (29). The latter are called *asymmetric* sums.

The following theorem due to Lam and Leung IV<sup>14</sup> will help us prove our theorem pertaining to vanishing sums of cosines (Theorem 2).

**Theorem 1 [Lam and Leung, Theorem 4.8].** *Let  $G$  be a cyclic group of order  $n = p_1^{a_1} p_2^{a_2} \cdots p_r^{a_r}$ , where  $p_1 < p_2 < \cdots < p_r$  are primes and let  $\varphi : \mathbb{Z}G \rightarrow \mathbb{Z}[\zeta]$  be as above, where  $\zeta = \zeta_n$ . For any minimal element  $x \in \mathbb{N}G \cap \ker(\varphi)$ , we have either (A)  $x$  is symmetric, or (B)  $r \geq 3$  and  $\epsilon_0(x) \geq p_1(p_2 - 1) + p_3 - p_2 > p_3$ .*

We shall utilize this theorem to prove the following (recall that  $n = N + 1$ ).

**Theorem 2.** *Let  $n$  be a positive odd integer and  $k_1, k_2, \dots, k_d$  be a set of integers such that  $1 \leq k_i \leq n - 1$ . Then*

$$\sum_{i=1}^d \cos\left(\frac{k_i \pi}{n}\right) \neq 0 \tag{30}$$

for any choice of  $k_i$ 's if and only if  $d$  is odd and is smaller than the smallest divisor of  $n$ .

*Proof.* By Lemma 3, we only need to consider  $d$  odd. Below we first work with roots of unity by writing the cosines in terms of the roots

$$\sum_{i=1}^d \cos\left(\frac{2k_i \pi}{2n}\right) = 2 \left\{ \sum_{i=1}^d \zeta_{2n}^{k_i} + \zeta_{2n}^{-k_i} \right\}. \tag{31}$$

So we have now a sum over  $2d$  roots of unity. We first prove that this sum is never zero if  $d < p_2$ . Since  $2n = 2p_2^{a_2} p_3^{a_3} \cdots p_r^{a_r}$  with all the  $p_i$ 's being odd, we are guaranteed (from Theorem IV) that  $p_1(p_2 - 1) + p_3 - p_2 = p_2 + p_3 - 2 < 2p_2$ ; therefore  $2d \equiv \epsilon_0(x) < 2p_2$  and if there were vanishing sums they would be of type (A), which are symmetric, i.e., sums of minimal relations. When  $d < p_2$ , in Eq. (31) there would be fewer than  $2p_2$  points on the unit circle, all of which appear as complex conjugate pairs. For the sum to be of type (A) and vanish, there should be a symmetric sum with a prime  $p$  that vanishes. The corresponding roots are a subset of the original points that are a vanishing sum of roots of  $\zeta_p$  with the prime  $p \geq p_2 > d$ ; therefore, it would involve a vanishing sum on more than half of the points of the original  $2d$  terms in Eq. (31). Hence there must be at least one complex conjugate pair in the vanishing sum under consideration. But if there is one complex conjugate pair, then all the roots should be complex conjugates as we can rotate any of the  $p$ th roots into one another.




FIG. 2. Left:  $\cos\left(\frac{k_1\pi}{N+1}\right) + \cos\left(\frac{k_2\pi}{N+1}\right) + \cos\left(\frac{k_3\pi}{N+1}\right) = 0$  is equivalent to the phasors adding to a vertical vector. The circles shown are unit circles. Right: Vanishing sums of roots of unity imply vanishing sums of cosines on the upper half plane since one can reflect any phasor without changing the cosine.

Since we have a vanishing sum of complex conjugate pairs but we allow only an odd number of terms, there must be a real root. But we exclude the real roots ( $\pm 1$ ). Therefore we reach a contradiction and the sum can never vanish.

Now we prove that the sum can be zero if  $d \geq p_2$ . It is sufficient to show that it vanishes for  $d = p_2$  as for any odd  $d > p_2$  we can always pair up the  $2(p_2 - d)$  cosines to cancel as we did in the proof of Lemma 3. Suppose  $d = p_2$ . Then a symmetric sum over the roots of unity that vanishes implies that the sum over cosines vanishes as the cosines are the real part and geometrically one can reflect the roots to the upper half plane (see Fig. 2). However, we need to exclude the possibility of  $\pm 1$  as roots and show that the sum still vanishes. The number of symmetric sums will be  $\frac{2n}{p}$  but only 2 of them have  $\pm 1$  as roots. In the sum involving the symmetric sums we can exclude the ones that have  $\pm 1$  and still be left with vanishing symmetric sums.  $\square$

*Corollary 2. The inverse of the Hückel matrix and hence its Green's function in  $d$  dimensions exists if and only if  $d$  is odd and is smaller than the smallest prime divisor of  $N + 1$ .*

For  $d = 3$ , this lemma and Eq. (30) have the geometrical interpretation shown in Fig. 2. Moreover, twice the left hand side of Eq. (30) is the expression for the energies of the Hückel matrix in  $d$  dimensions.

## V. THE PHYSICAL CONSEQUENCES OF THE INVERSE OF THE HÜCKEL MATRIX AND THE ZEROS OF ITS GREEN'S FUNCTION

The Hückel formalism, in its physical and chemical context, is not, of course, restricted to a linear chain. Various two- and three-dimensional connectivities have been probed in the 80 years of its existence, to the immense benefit of practice and understanding in chemistry. But until recent time, there has been scant interest in the Green's function of the Hückel matrix and its inverse. Heilbronner used the inverse of the Hückel Matrix to form an undervalued bridge between the resonance structure of valence bond theory and molecular orbitals—thus bringing together two seemingly distinct, but in fact related, approaches to the electronic structure of molecules.<sup>31</sup> The graph theoretical context has led people to investigate the inverse of the vertex adjacency matrix.<sup>10</sup> In the work of Estrada, the relationship between the Green's function formalism and the inverse of the vertex adjacency matrix of a graph is consistently utilized.<sup>8,9</sup>

In a field that has attracted much attention both experimentally and theoretically in the last decade, the transmission of current across molecules, a striking phenomenon, quite nonclassical, is observed. This is quantum interference, zero or low conductance when electrodes are attached to specific sites across a molecule.<sup>5,21</sup> Quantum interference occurs when the Green's function, whose absolute value squared is related to the current transmitted, vanishes. These are exactly the zeroes of

Eq. (12). The inverse of the Hückel matrix has been directly related to this phenomenon in the work of Markussen and Stadler.<sup>17</sup> The chemical consequences of just these zeroes have been outlined in recent work by us.<sup>24</sup>

The results we have obtained in this paper for the specific entries of the Green's function have been proven of great utility in describing the transmission of current through molecules. The Green's function expressions obtained in this paper also play an important role in designing new molecular switches based on electrocyclic reactions, cycloadditions, and sigmatropic reactions in linear polyenes.<sup>21,24</sup> In particular in a paper on linear polyenes, we have used the results to derive specifically the transmission across a chain and its exponential falloff.<sup>26</sup> In the same paper, where it was important to have the Green's function elements for a cyclic polyene (annulene) with and without bond alternation, expressions from the current paper and some based on similar procedures were used. Two kinds of zero values of the Green's function introduced in this paper, namely, easy zero and hard zero, provide the bases for an important classification of quantum interference, and the connection between these zeroes and the non-disjoint or disjoint nature of diradical molecules has been clarified.<sup>25</sup>

## ACKNOWLEDGMENTS

We thank Peter Shor for his help in proving Theorem 2. We also thank Christopher King, Jeffrey Goldstone, and Shev MacNamara. R.M. and R.H. thank National Science Foundation through Grant Nos. DMS 1312831 and CHE 1305872, respectively, G.S. is grateful for the ongoing support of MathWorks, and Y.T. thanks the Japan Society for the Promotion of Science for a JSPS Postdoctoral Fellowship for Research Abroad. R.M. thanks IBM Research for the support and the freedom provided by the Herman Goldstine fellowship.

## APPENDIX: FIRST PRINCIPLE DERIVATION FROM TRIGONOMETRICS

*Proposition.* The following sum has a closed form solution

$$-\frac{1}{N+1} \sum_{k=1}^N \frac{\sin(rk\omega) \sin(sk\omega)}{\cos(k\omega)} = e^{i\pi(\frac{r+s-1}{2})} \quad (\text{A1})$$

when  $r$  is even and  $s < r$  is odd. Otherwise, when  $s \leq r$  it is zero. Moreover  $s > r$  are symmetric, i.e.,  $G(r, s) = G(s, r)$ .

### 1. Easy zeros: Same parity of $s$ and $r$

Using  $\sin(rk\omega) \sin(sk\omega) = \frac{1}{2} \{ \cos[(r-s)k\omega] - \cos[(r+s)k\omega] \}$ , we can rewrite Eq. (7) as

$$G(r, s) = -\frac{1}{2(N+1)} \sum_{k=1}^N \frac{\{ \cos[(r-s)k\omega] - \cos[(r+s)k\omega] \}}{\cos(k\omega)}. \quad (\text{A2})$$

First let  $r-s = 2q$ ; this implies that  $r$  and  $s$  have the same parity (i.e., oddness or evenness). Therefore  $r+s$  is also even, and let it be  $r+s = 2q'$  for some  $q' \in \mathbb{N}$ . Eq. (A2) becomes

$$G(r, s) = -\frac{1}{2(N+1)} \sum_{k=1}^N \left\{ \frac{\cos[2qk\omega]}{\cos(k\omega)} - \frac{\cos[2q'k\omega]}{\cos(k\omega)} \right\}. \quad (\text{A3})$$

We now show that each sum is zero. Let us first show  $\sum_{k=1}^N \frac{\cos[2qk\omega]}{\cos(k\omega)} = 0$ . Recall  $\omega = \frac{\pi}{N+1}$  and expand the sum by adding the first to the last then the second to  $N-1$ st, etc., to get

$$\begin{aligned} \sum_{k=1}^N \frac{\cos[2qk\omega]}{\cos(k\omega)} &= \left( \frac{\cos[2q\omega]}{\cos(\omega)} + \frac{\cos[2qN\omega]}{\cos(N\omega)} \right) + \left( \frac{\cos[4q\omega]}{\cos(2\omega)} + \frac{\cos[2q(N-1)\omega]}{\cos((N-1)\omega)} \right) \\ &+ \dots + \left( \frac{\cos\left[2q\frac{N}{2}\omega\right]}{\cos(\omega N/2)} + \frac{\cos\left[2q\left(\frac{N}{2}+1\right)\omega\right]}{\cos\left(\left(\frac{N}{2}+1\right)\omega\right)} \right). \end{aligned} \quad (\text{A4})$$

We now show that each of the parenthesis is identically zero. To do so we notice that each of the parenthesis is of the form

$$\frac{\cos [2qk\omega]}{\cos (k\omega)} + \frac{\cos [2q(N-k+1)\omega]}{\cos ((N-k+1)\omega)}, \quad k = 1, 2, \dots, \frac{N}{2}.$$

But  $\cos((N-k+1)\frac{\pi}{N+1}) = \cos(-k\frac{\pi}{N+1} + \pi) = -\cos(k\frac{\pi}{N+1}) = -\cos(k\omega)$  by the double angle formula and evenness of the cosine. Moreover

$$\begin{aligned} \cos [2q(N-k+1)\omega] &= \cos \left[ 2q\pi - 2q\frac{k\pi}{N+1} \right] = \cos \left[ -2q\frac{k\pi}{N+1} \right] \\ &= \cos \left[ 2q\frac{k\pi}{N+1} \right] = \cos (2qk\omega). \end{aligned}$$

Concluding that the numerators are equal but denominators differ in sign resulting in

$$\frac{\cos [2qk\omega]}{\cos (k\omega)} + \frac{\cos [2q(N-k+1)\omega]}{\cos ((N-k+1)\omega)} = \frac{\cos [2qk\omega]}{\cos (k\omega)} - \frac{\cos [2qk\omega]}{\cos (k\omega)} = 0.$$

The exact same argument with substitution  $q'$  for  $q$  in Eq. (A4) proves that the second sum in Eq. (A3) is zero. Together proving  $G(r, s) = 0$  if  $r$  and  $s$  have the same parity. There are other zeros that are harder to prove.

**2. Harder zeros**

Let us make the sum in Eq. (7) centered by letting  $m = k - \frac{N+1}{2}$ , whereby

$$G(r, s) = -\frac{1}{N+1} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} \frac{\sin \left[ r\omega \left( m + \frac{N+1}{2} \right) \right] \sin \left[ s\omega \left( m + \frac{N+1}{2} \right) \right]}{\cos \left[ \omega \left( m + \frac{N+1}{2} \right) \right]},$$

where  $\omega = \frac{\pi}{N+1}$  as before. Since  $\cos[\omega(m + \frac{N+1}{2})] = -\sin(\omega m)$  and  $\sin x = \frac{1}{2i}(e^{ix} - e^{-ix})$ ,

$$\begin{aligned} G(r, s) &= -\frac{ie^{i(r+s)\pi/2}}{2(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} e^{i\omega m(r+s-1)} \frac{\left(1 - e^{-2ir\omega(m+\frac{N+1}{2})}\right) \left(1 - e^{-2is\omega(m+\frac{N+1}{2})}\right)}{1 - e^{-2i\omega m}} \\ &= -\frac{ie^{i(r+s)\pi/2}}{2(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} e^{i\omega m(r+s-1)} \frac{\left[1 - (-1)^r e^{-2ir\omega m}\right] \left[1 - (-1)^s e^{-2is\omega m}\right]}{1 - e^{-2i\omega m}}. \end{aligned} \tag{A5}$$

This equation is general and will be used later for nonzero sums as well.

Since we proved that if  $r$  and  $s$  have the same parity the sum vanishes, we prove the harder zeros (see Eq. (9)) by letting  $r$  be odd and  $s$  even and enforcing  $s < r$ . We can let  $r = 2q - 1$  and  $s = 2p$  with integers  $p$  and  $q$  satisfying  $0 < p < q \leq N/2$ . Using these, Eq. (A5) becomes

$$G(r, s) = -\frac{e^{i(p+q)\pi}}{2(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} e^{i2\omega m(p+q-1)} \frac{\left[1 + e^{-2i(2q-1)\omega m}\right] \left[1 - e^{-2i(2p)\omega m}\right]}{1 - e^{-2i\omega m}}.$$

We now use the factorization  $\frac{1-x^{2\ell}}{1-x} = 1 + x + x^2 + \dots + x^{2\ell-1}$  with  $x \equiv \exp(-i2m\omega)$  to get rid of the denominator

$$\begin{aligned} G(r, s) &= -\frac{(-1)^{p+q}}{2(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} \left\{ e^{i2\omega m(p+q-1)} \left[1 + e^{-2i\omega m(2q-1)}\right] \right. \\ &\quad \left. \times \left[1 + e^{-2i\omega m} + e^{-4i\omega m} + \dots + e^{-2i\omega m(2p-1)}\right] \right\}. \end{aligned}$$



Multiplying the phase factor into the parenthesis inside the sum and substituting for  $\omega$ , we have

$$G(r, s) = -\frac{(-1)^{p+q}}{2(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} \left( e^{2i(p+q-1)\frac{\pi m}{N+1}} + e^{-2i(q-p)\frac{\pi m}{N+1}} \right) \times \left\{ 1 + e^{-2i\frac{\pi m}{N+1}} + e^{-4i\frac{\pi m}{N+1}} + \dots + e^{-2i(2p-1)\frac{\pi m}{N+1}} \right\}. \tag{A6}$$

Comment: The pre-factor multiplying the sum can only be  $\frac{\pm 1}{2(N+1)}$ , determined by the values of  $p$  and  $q$ :  $G(r, s)$  vanishes iff the sum does.

We expand the summand in Eq. (A6) to get

$$\begin{aligned} G(r, s) &= -\frac{(-1)^{p+q}}{2(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} \left\{ \left[ e^{2i(p+q-1)\frac{\pi m}{N+1}} + e^{2i(p+q-2)\frac{\pi m}{N+1}} + \dots + e^{2i(q-p+1)\frac{\pi m}{N+1}} + e^{2i(q-p)\frac{\pi m}{N+1}} \right] \right. \\ &\quad \left. + \left[ e^{-2i(q-p)\frac{\pi m}{N+1}} + e^{-2i(q-p+1)\frac{\pi m}{N+1}} + e^{-2i(q-p+2)\frac{\pi m}{N+1}} + \dots + e^{-2i(p+q-2)\frac{\pi m}{N+1}} + e^{-2i(p+q-1)\frac{\pi m}{N+1}} \right] \right\} \\ &= -\frac{(-1)^{p+q}}{(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} \left\{ \cos \left[ \frac{2\pi m(q-p)}{N+1} \right] + \cos \left[ \frac{2\pi m(q-p+1)}{N+1} \right] + \dots \right. \\ &\quad \left. + \cos \left[ \frac{2\pi m(p+q-1)}{N+1} \right] \right\}, \end{aligned} \tag{A7}$$

where in the last equation, to get the cosines, we paired the first term inside the first brackets with the last term inside the second brackets etc. and used the formula  $e^{ix} + e^{-ix} = 2 \cos x$ . The factor of 2 cancelled the overall pre-factor 1/2.

Comment: It is important to note that, since  $q > p$ , the exponents in the first bracket are all positive and in the second bracket the exponents are all negative.

We can write a more succinct expression

$$G(r, s) = -\frac{(-1)^{p+q}}{(N+1)} \sum_{t=0}^{2p-1} \left\{ 2 \sum_{m=\frac{1}{2}}^{\frac{N-1}{2}} \cos \left[ \frac{2\pi m(q-p+t)}{N+1} \right] \right\}, \tag{A8}$$

where we used evenness of cosines, to let  $m$  run from 1/2, and switched the order of the sums. We now prove that the sum inside braces is  $(-1)^t$ . Let  $\theta = \frac{2\pi(q-p+t)}{N+1}$ ,  $n = m - 1/2$  and  $N' = \frac{N}{2} - 1$  to rewrite the sum

$$2 \sum_{m=\frac{1}{2}}^{\frac{N-1}{2}} \cos \left[ \frac{2\pi m(q-p+t)}{N+1} \right] \equiv 2 \sum_{n=0}^{N'} \cos \left[ \left( n + \frac{1}{2} \right) \theta \right]$$

but  $\cos \left[ \left( n + \frac{1}{2} \right) \theta \right] = \cos(n\theta) \cos\left(\frac{\theta}{2}\right) - \sin(n\theta) \sin\left(\frac{\theta}{2}\right)$  and Ref. 1,

$$\sum_{n=0}^{N'} \cos(n\theta) = \frac{\cos\left(\frac{N'\theta}{2}\right) \sin\left[\frac{\theta}{2}(N'+1)\right]}{\sin(\theta/2)}, \tag{A9}$$

$$\sum_{n=0}^{N'} \sin(n\theta) = \frac{\sin\left(\frac{N'\theta}{2}\right) \sin\left[\frac{\theta}{2}(N'+1)\right]}{\sin(\theta/2)}. \tag{A10}$$

Therefore  $\sum_{n=1}^{N'} \cos\left(n + \frac{1}{2}\right)\theta = \sum_{n=1}^{N'} \left\{ \cos(n\theta) \cos\left(\frac{\theta}{2}\right) - \sin(n\theta) \sin\left(\frac{\theta}{2}\right) \right\}$  gives

$$\begin{aligned} 2 \sum_{n=0}^{N'} \cos\left(n + \frac{1}{2}\right)\theta &= 2 \left\{ \cos\left(\frac{\theta}{2}\right) \frac{\cos\left(\frac{N'\theta}{2}\right) \sin\left[\frac{\theta}{2}(N'+1)\right]}{\sin(\theta/2)} - \sin\left(\frac{\theta}{2}\right) \frac{\sin\left(\frac{N'\theta}{2}\right) \sin\left[\frac{\theta}{2}(N'+1)\right]}{\sin(\theta/2)} \right\} \\ &= 2 \frac{\sin\left[\frac{\theta}{2}(N'+1)\right]}{\sin(\theta/2)} \left\{ \cos\left(\frac{N'\theta}{2}\right) \cos\left(\frac{\theta}{2}\right) - \sin\left(\frac{N'\theta}{2}\right) \sin\left(\frac{\theta}{2}\right) \right\} \\ &= 2 \frac{\sin\left[\frac{\theta}{2}(N'+1)\right]}{\sin(\theta/2)} \left\{ \cos\left(\frac{\theta}{2}(N'+1)\right) \right\} = \frac{\sin[(N'+1)\theta]}{\sin(\theta/2)} = \frac{\sin[N\theta/2]}{\sin(\theta/2)}. \end{aligned}$$

However  $\sin[N\theta/2] = \sin\left[\frac{(N+1)\theta}{2} - \frac{\theta}{2}\right] = \sin\left(\frac{(N+1)\theta}{2}\right) \cos\frac{\theta}{2} - \cos\left(\frac{(N+1)\theta}{2}\right) \sin\frac{\theta}{2}$ . But  $\sin\left(\frac{(N+1)\theta}{2}\right) = 0$ , leaving us with

$$\frac{\sin[N\theta/2]}{\sin(\theta/2)} = \frac{\sin\left[\frac{\pi N(q-p+t)}{N+1}\right]}{\sin\left(\frac{\pi(q-p+t)}{N+1}\right)} = -\cos\left(\frac{(N+1)\theta}{2}\right) = -\cos(\pi(q-p+t)) = -(-1)^{q-p+t}. \tag{A11}$$

Putting this back into the sum (Eq. (A8))

$$\begin{aligned} G(r, s) &= \frac{(-1)^{p+q}}{(N+1)} \sum_{t=0}^{2p-1} \{(-1)^{q-p+t}\} \\ &= \frac{(-1)^{2q}}{(N+1)} \sum_{t=0}^{2p-1} (-1)^t = \frac{1}{(N+1)} \sum_{t=0}^{2p-1} (-1)^t, \end{aligned}$$

zero comes out because we are summing alternating +1's and -1's an even number of times. This completes the proof of the harder zeros. Note that we used  $q > p$ . For example if  $q = p$ , then  $\cos \pi(q - p + t)$  would be 1 for  $t = 0$  and the sum would give a  $2p - 1$  on that term alone.

Recall  $\frac{r+1}{2} = q$  and  $\frac{s}{2} = p$  with integers  $p$  and  $q$  satisfying  $0 < p < q \leq N/2$ ; for this choice

$$G(r, s) = 0.$$

### 3. Nonzero entries: ±1's in the G

It remains to show that when  $r$  is even and  $s$  is odd,  $G(r, s)$  is  $\pm 1$  as shown in Eq. (9). Let  $r = 2q$  and  $s = 2p - 1$  with  $p \leq q$  (note that we allow for equality as well). Using Eq. (A5) and previous techniques, we have

$$\begin{aligned} G(r, s) &= -\frac{ie^{i(2(p+q)-1)\pi/2}}{2(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} e^{2i\omega m(p+q-1)} \frac{[1 - e^{-2i(2q)\omega m}][1 + e^{-2i(2p-1)\omega m}]}{1 - e^{-2i\omega m}} \\ &= -\frac{ie^{i(2(p+q)-1)\pi/2}}{2(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} (e^{2i\omega m(p+q-1)} + e^{-2i\omega m(p-q)}) \\ &\quad \times \{1 + e^{-2i\omega m} + e^{-4i\omega m} + \dots + e^{-2i(2q-1)\omega m}\}. \end{aligned}$$

Once again we multiply the parenthesis into the braces to get (using  $ie^{i(2(p+q)-1)\pi/2} = e^{i(p+q)\pi}$ ),

$$\begin{aligned} G(r, s) &= -\frac{e^{i(p+q)\pi}}{2(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} \left\{ [e^{2i\omega m(p+q-1)} + e^{2i\omega m(p+q-2)} + \dots + e^{2i\omega m(p-q)}] \right. \\ &\quad \left. + [e^{-2i\omega m(p-q)} + e^{-2i\omega m(p-q+1)} + \dots + e^{-2i\omega m(p+q-1)}] \right\}. \end{aligned} \tag{A12}$$

Comment: Eq. (A12) looks very similar to Eq. (A7); however, it has a key difference. Since  $q \geq p$ , in either one of the brackets there will be a term with exponent zero. For example, if one looks

at the first brackets the first term is  $e^{2i\omega m(p+q-1)}$ , which clearly has a positive exponent; however, the last term  $e^{2i\omega m(p-q)}$  has either zero or negative exponent. If it is negative, then a term preceding it must have had zero exponent. Therefore, the sum for some choice of  $r$  and  $s$  can look like

$$G_{\text{example}}(r, s) = -\frac{e^{i(p+q)\pi}}{2(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} \left\{ \left[ e^{2i\omega m(p+q-1)} + \dots + e^{2i\omega m} + 1 + e^{-2i\omega m} + \dots + e^{2i\omega m(p-q)} \right] \right. \\ \left. + \left[ e^{-2i\omega m(p-q)} + e^{2i\omega m} + 1 + e^{-2i\omega m} + \dots + e^{-2i\omega m(p+q-1)} \right] \right\}. \tag{A13}$$

We can pair the terms to the left (right) of the 1 in the first bracket with those to the right (left) of the 1 in the right bracket to get the cosines as before. It is clear that the sum over 2 contributes a  $2(N-1)$ . We now show that the sum over the cosines contributes a 4, which together makes  $2(N+1)$  and cancels the denominator in the pre-factor.

For any  $p$  and  $q$ , we can find a  $t_0 = q - p \geq 0$  that makes the exponent zero. In the first bracket, there are  $q - p$  terms to its left and there are  $2q - (q - p + 1) = p + q - 1$  terms to its right (for a total of  $2q$  terms). We can pair the terms to its left with the corresponding terms in the second bracket (now to the right of the 1) to get cosines and similarly pair terms to its right to get cosines. Then, we can break the sum in the foregoing equation to the sum over cosines obtained from terms to the left of  $t_0$  in the first bracket, the sum over terms to its right and add a 2 for the term itself. Namely,

$$G(r, s) = -\frac{e^{i(p+q)\pi}}{2(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} \left\{ 2 \sum_{t=1}^{q-p} \cos [2\omega mt] + 2 + 2 \sum_{t=1}^{p+q-1} \cos [2\omega mt] \right\} \tag{A14}$$

$$= -\frac{e^{i(p+q)\pi}}{(N+1)} \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} \left\{ \sum_{t=1}^{q-p} \cos [2\omega mt] + \sum_{t=1}^{p+q-1} \cos [2\omega mt] + 1 \right\},$$

where we cancelled the overall pre-factor of a  $1/2$ . Let us evaluate each of the sums separately (switching order of summation, changing variables as before)

$$\sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} \cos [2\omega mt] = 2 \sum_{n=0}^{\frac{N}{2}-1} \cos \left[ 2\omega t \left( n + \frac{1}{2} \right) \right]$$

$$= 2 \cos(\omega t) \sum_{n=0}^{\frac{N}{2}-1} \cos(2\omega tn) - 2 \sin(\omega t) \sum_{n=0}^{\frac{N}{2}-1} \sin(2\omega tn).$$

The sum over cosines is evaluated using Eqs. (A9) and (A10),

$$\sum_{n=0}^{\frac{N}{2}-1} \cos(2\omega tn) = \frac{\cos \left( \left( \frac{N}{2} - 1 \right) \omega t \right) \sin [\omega t (N/2)]}{\sin(\omega t)}$$

$$\sum_{n=0}^{\frac{N}{2}-1} \sin(2\omega tn) = \frac{\sin \left( \left( \frac{N}{2} - 1 \right) \omega t \right) \sin [\omega t (N/2)]}{\sin(\omega t)},$$

which together give

$$\sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} \cos [2\omega mt] = 2 \frac{\sin \left[ \left( \frac{N}{2} \right) \omega t \right]}{\sin(\omega t)} \left\{ \cos \left( \left( \frac{N}{2} - 1 \right) \omega t \right) \cos(\omega t) - \sin \left( \left( \frac{N}{2} - 1 \right) \omega t \right) \sin(\omega t) \right\}$$

$$= 2 \frac{\sin \left[ \left( \frac{N}{2} \right) \omega t \right]}{\sin(\omega t)} \left\{ \cos \left[ \left( \frac{N}{2} \right) \omega t \right] \right\} = \frac{\sin(N\omega t)}{\sin(\omega t)} = \frac{\sin \left( \frac{\pi Nt}{N+1} \right)}{\sin \left( \frac{\pi t}{N+1} \right)}.$$

We calculated this ratio in Eq. (A11) so we have

$$\frac{\sin\left(\frac{\pi Nt}{N+1}\right)}{\sin\left(\frac{\pi t}{N+1}\right)} = -(-1)^t \quad .$$

Using this we can evaluate Eq. (A14),

$$G(r, s) = -\frac{e^{i(p+q)\pi}}{(N+1)} \left\{ -\sum_{t=1}^{q-p} (-1)^t - \sum_{t=1}^{p+q-1} (-1)^t + \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} 1 \right\}.$$

If  $q-p$  is even then  $q-p = 2k$  for some  $k$  and  $p+q-1 = 2(k+p) - 1$ , which is odd. Also if  $q-p = 2k-1$ , then  $p+q-1 = 2(p+k-1)$ , which is even. In either case one of the sums vanishes and the other evaluates to be  $-1$ . Therefore,

$$G(r, s) = -\frac{e^{i(p+q)\pi}}{(N+1)} \left\{ 1 + \sum_{m=-\frac{N-1}{2}}^{\frac{N-1}{2}} 1 \right\} = -\frac{e^{i(p+q)\pi}}{(N+1)} (N+1) = -e^{i(p+q)\pi}. \quad (\text{A15})$$

So we predict that if  $p+q$  is odd then  $G(r, s) = +1$  and if  $p+q$  is even then  $G(r, s) = -1$ . What does this mean for  $r$  and  $s$ ? Let us cover all of the cases one by one. Recall that  $r$  is even and  $s$  is odd and  $q+p = \frac{r}{2} + \frac{s+1}{2}$ .

- $p+q$  is even,  $G(r, s) = -1$ , and  $q$  is even. This means that  $p$  is even. These imply that  $r$  and  $s+1$  are multiples of 4. Looking at  $\mathbf{G}$ , we see that these entries indeed are  $-1$ .
- $p+q$  is even,  $G(r, s) = -1$ , and  $q$  is odd. This means  $p$  is odd. These imply that  $r$  and  $s+1$  are multiples of 2 but not 4. Looking at  $\mathbf{G}$ , we see that the rest of the entries that are  $-1$  have been covered.
- $p+q$  is odd,  $G(r, s) = +1$ , and  $q$  is even. This means  $p$  is odd. These imply that  $r$  is a multiple of 4 but  $s+1$  is not (though of course even). This covers some of the  $+1$ 's in  $\mathbf{G}$ .
- Lastly,  $p+q$  is odd,  $G(r, s) = +1$ , and  $q$  is odd. This means  $p$  is even. These imply that  $r$  is not a multiple of 4 (though of course even), yet  $s+1$  is a multiple of 4. These cover the rest of  $+1$ 's seen  $\mathbf{G}$ .

The final result Eq. (A15) can be expressed in terms of  $r$  and  $s$  as

$$G(r, s) = -e^{i\pi\left(\frac{r+s+1}{2}\right)} = e^{i\pi\left(\frac{r+s-1}{2}\right)}. \quad (\text{A16})$$

This completes our proof.

*Remark 4.* All the equations above for  $G(r, s)$  were checked numerically.

<sup>1</sup> Wolfram MathWorld (search under "cosine" and "sine").

<sup>2</sup> S. Alvarez, *Tables of Parameters for Extended Hückel Calculations* (Universitat de Barcelona, Barcelona, 1989).

<sup>3</sup> N. W. Ashcroft and N. David Mermin, *Solid State Physics* (Saunders College, Philadelphia, 1976 and 1981).

<sup>4</sup> C. A. Coulson and G. S. Rushbrooke, "Note on the method of molecular orbitals," in *Mathematical Proceedings of the Cambridge Philosophical Society* (Cambridge University Press, 1940), Vol. 36, pp. 193–200.

<sup>5</sup> S. Datta, *Quantum Transport: Atom to Transistor* (Cambridge University Press, Cambridge, 2005).

<sup>6</sup> P. J. Davis, *Circulant Matrices* (American Mathematical Society, 1979), pp. 66–101.

<sup>7</sup> P. A. M. Dirac, *The Principles of Quantum Mechanics*, 4th ed. (Oxford University Press, 1967).

<sup>8</sup> E. Estrada and N. Hatano, "Statistical-mechanical approach to subgraph centrality in complex networks," *Chem. Phys. Lett.* **439**(1), 247–251 (2007).

<sup>9</sup> E. Estrada and N. Hatano, "Communicability in complex networks," *Phys. Rev. E* **77**, 036111 (2008).

<sup>10</sup> A. Farrugia, J. B. Gauci, and I. Sciriha, "On the inverse of the adjacency matrix of a graph," *Spec. Matrices* **1**, 28–41 (2013).

<sup>11</sup> R. M. Gray, "Toeplitz and circulant matrices: A review," *Commun. Inf. Theory* **2**(3), 155–239 (2005).

<sup>12</sup> I. Gutman and O. E. Polonsky, *Mathematical Concepts in Organic Chemistry* (Springer, Berlin, 1986).

<sup>13</sup> E. Heilbronner and H. Bock, *The HMO-Model and Its Application* (Wiley, London, 1976).

<sup>14</sup> T. Y. Lam and K. H. Leung, "On vanishing sums of roots of unity," *J. Algebra* **224**, 91–109 (2000).

<sup>15</sup> H. C. Longuet-Higgins and L. Salem, "The alternation of bond lengths in large conjugated molecules. III. The cyclic polyenes  $C_{18}H_{18}$ ,  $C_{24}H_{24}$ , and  $C_{30}H_{30}$ ," *Proc. R. Soc. A* **257**, 445–456 (1960).

<sup>16</sup> R. K. Mallik, "The inverse of a tridiagonal matrix," *Linear Algebra Appl.* **325**(1), 109–139 (2001).

- <sup>17</sup> T. Markussen, R. Stadler, and K. S. Thygesen, "Graphical prediction of quantum interference-induced transmission nodes in functionalized organic molecules," *Phys. Chem. Chem. Phys.* **13**, 14311–14317 (2011).
- <sup>18</sup> G. Meurant, "A review on the inverse of symmetric tridiagonal and block tridiagonal matrices," *SIAM J. Matrix Anal. Appl.* **13**(3), 707–728 (1992).
- <sup>19</sup> B. Poonen and M. Rubinstein, "The number of intersection points made by the diagonals of a regular polygon," *SIAM J. Discrete Math.* **11**, 135–156 (1998).
- <sup>20</sup> P. Schlegel, "The explicit inverse of a tridiagonal matrix," *Math. Comput.* **24**(111), 665 (1970).
- <sup>21</sup> G. C. Solomon, C. Herrmann, and M. Ratner, "Molecular electronic junction transport: Some pathways and some ideas," *Top. Curr. Chem.* **313**, 1 (2012).
- <sup>22</sup> G. Strang, "The discrete cosine transform," *SIAM Rev.* **41**(1), 135–147 (1999).
- <sup>23</sup> L. N. Trefethen and M. Embree, *Spectra and Pseudospectra* (Princeton University Press, 2005).
- <sup>24</sup> Y. Tsuji, R. Hoffmann, R. Movassagh, and S. Datta, "Quantum interference in polyenes," *J. Chem. Phys.* **141**(22), 224311 (2014).
- <sup>25</sup> Y. Tsuji, R. Hoffmann, M. Strange, and G. C. Solomon, "Close relation between quantum interference in molecular conductance and diradical existence," *Proc. Natl. Acad. Sci. U. S. A.* **113**(4), E413–E419 (2016).
- <sup>26</sup> Y. Tsuji, R. Movassagh, S. Datta, and R. Hoffmann, "Exponential attenuation of through-bond transmission in a polyene: Theory and potential realizations," *ACS Nano* **9**(11), 11109–11120 (2015).
- <sup>27</sup> R. A. Usmani, "Inversion of a tridiagonal Jacobi matrix," *Linear Algebra Appl.* **212-213**, 413–414 (1994).
- <sup>28</sup> R. Vandebril, M. Van Barel, and N. Mastronardi, *Matrix Computations and Semiseparable Matrices: Linear Systems* (JHU Press, 2007), Vol. 1.
- <sup>29</sup> C. S. Wannere, K. W. Sattelmeyer, H. F. Schaefer, and P. von Ragué Schleyer, "Aromaticity: The alternating C–C bond length structures of [14]-, [18]-, and [22] annulene," *Angew. Chem.* **116**(32), 4296–4302 (2004).
- <sup>30</sup> Y. Xue, S. Datta, and M. A. Ratner, "First-principles based matrix Green's function approach to molecular electronic devices: General formalism," *Chem. Phys.* **281**(2), 151–170 (2002).
- <sup>31</sup> E. Heilbronner, "Über einen graphentheoretischen Zusammenhang zwischen dem Hückel'schen MO-Verfahren und dem Formalismus der Resonanztheorie," *Helv. Chim. Acta* **45**, 1722–1725 (1962).